

# INVENTORY AND DIVERSITY OF BOTANICAL PESTICIDES: A CASE STUDY OF THE BLIDIAN ATLAS, NORTH-CENTRAL ALGERIA

BOURENANE, E.<sup>1\*</sup> – MOHAMMEDI, A.<sup>1</sup> – AMOKRANE, D.<sup>1,2</sup> – MEKIDECHE, S.<sup>3</sup> – MEZANI, S.<sup>4</sup> – ABABOU, A.<sup>5</sup>

<sup>1</sup>*Laboratory of Local Natural Bio-Resources (LBRN), Department of Agronomy, Faculty of Nature and Life Sciences, University Hassiba Ben Bouali of Chlef, Chlef 02000, Algeria (phone: +213-027-7270)*

<sup>2</sup>*Laboratory of Crop Production and Protection in the region of Chlef (PPCRC), Department of Agronomy, Faculty of Nature and Life Sciences, University Hassiba Ben Bouali of Chlef, Chlef 02000, Algeria*

<sup>3</sup>*Laboratory of Plant ecology and environment ((LEVE), Department of Ecology and Environment, Faculty of Biological Sciences, University of Sciences and technology, Houari Boumediene, Algiers 16000, Algeria (phone: +213-021-247-311)*

<sup>4</sup>*Laboratory of Natural Ressources (LRN), Department of Ecology and Environment, Faculty of Biological Sciences and Agronomic Sciences, University Mouloud Mammeri of Tizi-Ouzou, Tizi-Ouzou 15000, Algeria (phone: +213-026-115-651)*

<sup>5</sup>*Laboratory of Local Natural Bio-Resources (LBRN), Department of Biology, Faculty of Nature and Life Sciences, University Hassiba Ben Bouali of Chlef, Chlef 02000, Algeria*

\*Corresponding author

e-mail: sbourenane313@gmail.com; phone: +213-549-066-829

(Received 1<sup>st</sup> Jan 2025; accepted 19<sup>th</sup> Mar 2025)

**Abstract.** Botanical pesticides are receiving increasing attention from researchers and growers as an alternative to synthetic pesticides. In this study, we identified the pesticidal plants naturally occurring in the Blidian Atlas in North Algeria during the flowering period of the most of plants (mid-March to mid-June). These plants were identified in different habitats: forest, wasteland and maquis across three vegetation zones: Thermo-Mediterranean, Meso-Mediterranean and Supra-Mediterranean. The study recorded notable floristic diversity. Among the 452 total species recorded in the area, 88 species distributed across 39 botanical families were effective against pest insects and fungi. Asteraceae and Lamiaceae were the most dominant families comprising 18.18% and 13.63%, respectively. In terms of biological types, hemicryptophyte and therophyte were the most dominant with 30.68% and 25%, respectively. The highest floristic diversity was recorded in the Meso-Mediterranean stage, particularly in the maquis and forest areas. Insecticidal compounds of most of the identified plants were found primarily in the leaves followed to a lesser extent, by the flowers, while the most targeted insects were storage insects.

**Keywords:** *plant pesticides, vegetation-zones, pests, fungi, habitats*

## Introduction

To meet the growing demand for staple agricultural products from the global population, the use of synthetic chemical pesticides has become a fast and efficient way to protect crops from pest infestations (Lengai et al., 2019), they act directly or

indirectly as vectors of plant diseases, contributing to a decline in both the quality and quantity of crops (Zettler et al., 1990).

However, the excessive and indiscriminate use of synthetic pesticides can have detrimental effects on human health (Sayed et al., 2022) and the environment (Sande et al., 2011) including soil and groundwater contamination, ozone layer depletion, and threats to biodiversity. This is primarily due to their non-biodegradable nature (Wimalawansa et al., 2014), toxicity to non-target organisms (Sande et al., 2011) and their ability to promote resistance in insects and pathogens when used over prolonged periods (Lengai et al., 2019).

It is therefore essential to explore safe alternatives to synthetic pesticides such as botanical pesticides, which are believed to be effective against a variety of crop pests (Acheuk et al., 2022). The use of botanical pesticides promotes a healthy environment and supports sustainable agriculture (Shabana et al., 2017). Their importance lies in several factors including their efficacy, biodegradability, and low toxicity. Additionally, they offer a wide range of modes of action and are easily accessible (Neeraj et al., 2017). Furthermore, botanical pesticides are a rich source of bioactive compounds (Qin et al., 2010) that serve as insecticides, repellents, bactericides, fungicides, herbicides, and nematicides (Isman et al., 2006).

In this regard, over 2000 plant species have been identified worldwide for their traditional use as insecticides (Adeniyi et al., 2010). The insecticidal toxicity of these plants has garnered significant attention from researchers in recent years, reflected in the growing number of publications focused on this subject.

Despite the presence of over 3000 spontaneous plant species in Algeria (Quezel and Santa, 1962-1963), studies on the inventory of botanical pesticides in the country remain limited. However, researchers are starting to recognize their significance (Amokrane et al., 2023), especially in mountainous regions. These areas are known for their rich biodiversity, closely associated with the region's topographic and bioclimatic variability (Mekideche et al., 2018).

In this context, the present study aims to create an inventory of pesticidal plants in the Blidian Atlas massif, a significant plant-rich region. It will also synthesize key research on these plants based on existing literature, covering their botanical families, biological types, habitats, phytogeographies, targeted pests and pathogens and the specific plant parts used for biological control. Additionally, the biological trials conducted by researchers on these plants will be reviewed.


This study will serve as a valuable database for future research and the use of bioinsecticides in Algeria and beyond.

## **Materials and methods**

### ***The study region***

The Blidean Atlas is located in northern Algeria, within the province of Blida, between 36°15'00" and 36°45'00" North and 2°30'00" and 3°50'00" East, covering an area of 52145 hectares (*Fig. 1*). It belongs to the Mediterranean climate, characterized by wet winters and hot, dry summers.

The analysis of climate data from 1999 to 2013 reveals a subhumid to per-humid bioclimate, with temperate winters. The average annual precipitation ranges from 679 to 1382 mm, while the mean annual temperatures vary between 11°C and 19.2°C (Mekideche et al., 2018).



**Figure 1.** Location of the study stations (Hammam Melouane, Aissaouia, Chrea) and the main vegetation zones (Thermo-mediterranean, Meso-mediterranean, Supra-mediterranean) in the Blidean Atlas (Algeria)

### Sampling and data analysis

The inventories were conducted over a period of two years (2023 and 2024) and include a total species of 452, involving a total of 48 random surveys in the three bioclimatic zones: Thermo-Mediterranean, Meso-Mediterranean, and Supra-Mediterranean spanning an altitudinal gradient from 250 to 1300 meters. Each bioclimatic zone was further divided into three habitats: wastelands, maquis, and forests.

According to Meddour (1994), the Thermo-Mediterranean zone is located below 600 m in altitude and is characterized by a subhumid bioclimatic stage, the vegetation dominated by *Pinus halepensis*, *Olea europaea*, and *Pistacia lentiscus*. The Meso-Mediterranean zone, situated between 600 and 1300 m, features humid bioclimatic stage dominated by *Quercus suber* and *Quercus ilex*. The Supra-Mediterranean zone, found above 1300 m, is characterized by perhumid bioclimatic stage, dominated by *Cedrus atlantica*.

Botanical identification was based on specimens from the herbarium of Chrea National Park. The family of each species was determined according to the APG IV (2016) classification, the biological types were distinguished according to Raunkiaer (1934), and the phytogeographical distribution according to Quézel and Santa (1962-1963).

The species diversity associated with each bioclimatic zone and habitat was calculated using Shannon diversity index ( $H'$ ), which combines species richness and evenness (Shannon and Weaver, 1949). It is calculated as follows:

$$H' = \sum_{i=1}^s pi \log_2 pi \quad (\text{Eq.1})$$

where:  $H'$  = The Shannon index value,  $p_i$  = the proportion of individuals belonging to the  $i$ th species,  $\log_2$  = the natural logarithm and  $S$  = the number of species in the community.

To study the relationship between the different species and their habitats, we used Correspondence Analysis (CA), a multivariate technique that examines associations and oppositions within the data, using the R software.

## Results

Among 452 species recorded in the study area, 88 were effective against pests and phytopathogenic agents (Table 1), distributed among 39 distinct botanical families. The Asteraceae family contained the largest number of species, with 16 species (18%), followed by the Lamiaceae with 12 species (14%), the Apiaceae with 8 species (9%), and the Euphorbiaceae with 4 species (5%). The Myrtaceae, Cistaceae, and Fabaceae families each had 3 species (3%). The Hypericaceae, Brassicaceae, Pinaceae, Plantaginaceae, Malvaceae, and Caprifoliaceae families with 2 species (2%), while the Oleaceae, Asphodelaceae, Urticaceae, Lauraceae, Anacardiaceae, Convolvulaceae, Orobanchaceae, Dennstaedtiaceae, Scrophulariaceae, Thymelaeaceae, Rosaceae, Resedaceae, Violaceae, Capparaceae, Caryophyllaceae, Linaceae, Verbenaceae, Taxaceae, Polygonaceae, Papaveraceae, Aristolochiaceae, Boraginaceae, Apocynaceae, and Oxalidaceae each with only one species (1%) (Fig. 2). The aerial parts, particularly the leaves, were the most commonly used for extracting bioinsecticidal molecules. Most of these biopesticides were applied against storage pests, such as *Tribolium castaneum*, *Sitophilus granarius*, *Tribolium confusum*, *Callosobruchus maculatus*, and *Rhyzopertha dominica* (Table 1).

In terms of biological types, hemicryptophytes (Hem) were the most represented, with 27 species (31%), followed by therophytes (The) and phanerophytes (Pha), with 22 species (25%) and 21 species (24%), respectively. chamaephytes (Cha) accounted for 14 species (16%), while geophytes (Geo) comprised only 4 species (4%) (Fig. 3).

Most of the inventoried species were of Mediterranean origin, with 43 species (49%), followed by cosmopolitan species with 9 species (10%), Euro-Asiatic species with 8 species (9%) and European species with 5 species (6%). 4 species (5%) were from Euro-Mediterranean origin, while Atlantico-Mediterranean and endemic plants each accounted for 3 species (3%). Australian, Mediterranean-Saharo-Sindian and Paleotemperate species each represented by 2 species (2%). Finally, Oro-Mediterranean, Ibero-Mauritian, subtropical and South African species accounted for 1 species each (1%) (Fig. 4).

The highest diversity was recorded in the Maottorals (ranging from 4.2 to 4.5), followed by wastelands (4.2 to 4.3), while forests were the least diverse habitats (3.2 to 3.7). In terms of vegetation zones, the meso-Mediterranean and supra-Mediterranean zones exhibited the highest diversity, with Shannon indices (Eq. 1) of 4.17 and 4.14, respectively, while the thermo-Mediterranean zone was less diverse, with a Shannon index of 3.87 (Table 2). Overall, the results indicate that the study area displays very high diversity.

**Table 1.** Inventoried pesticide plants, used parts and target pests

Species	Pest	Plant part	References
<i>Anacyclus clavatus</i> (Desf.) Pers.	<i>Tribolium castaneum</i> (H) Herbst	LE	Pascual-Villalobos et al. (1998)
<i>Atractylis gummifera</i> (L)	<i>Bactrocera oleae</i> (Rossi.)	NM	Mejdoub et al. (2020)
	<i>Sitophilus granarius</i> (L)	RO	Chelef et al. (2021)
<i>Atractylis cancellata</i> (L)	<i>Tribolium castaneum</i> (H).Herbst	WP	Pascual-Villalobos et al. (1998)
<i>Calendula arvensis</i> Boiss.	<i>Tribolium castaneum</i> (H) Herbst	WP	Pascual-Villalobos et al. (1998)
<i>Carthamus lanatus</i> (L)	<i>Spodoptera littoralis</i> (Bois.)	ST	Pavela (2011)
<i>Coleostephus myconis</i> (L)	<i>Tribolium confusum</i> (J. Du Val)	FL/LE	Haouas et al. (2014)
<i>Dittrichia viscosa</i> (L)	<i>Callosobruchus aculatus</i> (Fab.)	AP	Gueribis et al. (2019)
	<i>Spodoptera littoralis</i> (Bois.) / <i>Myzus persicae</i> (Sulzer) / <i>Rhopalosiphum padi</i> (Linnaeus)	AP	Mamoci et al. (2012)
	<i>Verticillium dahliae</i> Kleb	NM	Varo et al. (2017)
<i>Echinops spinosissimus</i> Turra	<i>Bactrocera oleae</i> (Rossi.)	NM	Belabbes et al. (2020)
<i>Erigeron canadensis</i> (L).	<i>Tribolium castaneum</i> (H) Herbst / <i>Aspergillus flavus</i> (L)	AP	Azeem et al. (2022)
<i>Glebionis coronaria</i> (L.) Cass. ex Spach	<i>Alternaria brassicicola</i> (Schwein.) / <i>Aspergillus flavus</i> (L)/ <i>Botrytis cinerea</i> Pers. / <i>Fusarium verticillioides</i> (Sacc.) / <i>Fusarium solani</i> (Mart.) Sacc. / <i>Mycocentrospora acerina</i> (R. Hartig) / <i>Pythium ultimum</i> Trow / <i>Rhizoctonia solani</i> Kühn / <i>Sclerotinia sclerotiorum</i> (Lib.) / <i>Serpula lacrymans</i> (Wulfen)	NM	Alvarez-Castellanos et al. (2001)
	<i>Tribolium castaneum</i> (H).Herbst. / <i>Rhyzopertha dominica</i> (Fab.)	NM	Hazafa et al. (2022)
<i>Helichrysum stoechas</i> (L.)	<i>Tribolium castaneum</i> (H).Herbst	NM	Pascual-Villalobos et al. (1998)
<i>Mauranthemum paludosum</i> (Poir.) Vogt & Oberpr.	<i>Tribolium confusum</i> (J. Du Val)	FL/LE	Haouas et al. (2014)
<i>Pallenis spinosa</i> (L.) Cass	<i>Tribolium castaneum</i> (H).Herbst.	WP	Pascual-Villalobos et al. (1998)
<i>Sonchus asper</i> (L.) Hill	<i>Tribolium castaneum</i> (H).Herbst.	RO/LE	Samreen et al. (2014)
	<i>Helicoverpa armigera</i> Hübner/ <i>Spodoptera litura</i> (Fab.)	NM	Shehzad et al. (2021)
<i>Sonchus oleraceus</i> (L.)	<i>Sitophilus oryzae</i> (L.)	LE	Shawer et al. (2020)
<i>Senecio vulgaris</i> (L.)	<i>Tribolium castaneum</i> (H).Herbst.	WP	Pascual-Villalobos et al. (1998)

Species	Pest	Plant part	References
<i>Ajuga iva</i> (L.) Schreb.	<i>Tribolium castaneum</i> (H) Herbst	NM	Jbilou et al. (2008)
	<i>Plodia interpunctella</i> (Hübner)	AP	Bouayad et al. (2013)
<i>Clinopodium nepeta</i> (L.) Kuntze	<i>Aspergillus flavus</i> (L)/ <i>Tribolium confusum</i> (J. Du Val)/ <i>Sitophilus zeamais</i> (L).	NM	Debbabi et al. (2020)
<i>Clinopodium vulgare</i> (L.)	<i>Spodoptera littoralis</i> (Bois.)	ST	Pavela (2011)
<i>Lavandula stoechas</i> (L.)	<i>Tetranychus cinnabarinus</i> (Bois.)	AP	Sertkaya et al. (2010)
	<i>Ceratitis capitata</i> (Wiedemann)	NM	Ghalbane et al. (2021)
	<i>Spodoptera littoralis</i> (Bois.)/ <i>Leptinotarsa decemlineata</i> (Say.)/ <i>Myzus persicae</i> (Sulzer)	FL/LE	Gonzalez-Coloma et al. (2006)
<i>Lavandula dentata</i> (L.)	<i>Sitophilus zeamais</i> / <i>Tribolium castaneum</i> (L)/ <i>Epicauta atomaria</i> (Germar)	ST/LE	Wagner et al. (2021)
	<i>Ceratitis capitata</i> (Wiedemann)	NM	Ghalbane et al. (2021)
<i>Mentha rotundifolia</i> (L.) Huds.	<i>Fusarium culmorum</i> / <i>Sitophilus granarius</i> (Linnaeus) / <i>Tribolium confusum</i> (J)	ST/LE	Yakhlef et al. (2020)
<i>Salvia verbenaca</i> (L.)	<i>Callosobruchus chinensis</i> (Linnaeus)	LE	Righi et al. (2014)
<i>Origanum floribundum</i> Munby	<i>Sitobion avenae</i> (F)	LE	Mouas et al (2021)
<i>Satureja montana</i> (L)	<i>Spodoptera littoralis</i> (Bois.)/ <i>Myzus persicae</i> (Sulzer) / <i>Leptinotarsa decemlineata</i> (Say.)	LE / FL	Navarro-Rocha et al. (2020)
	<i>Acyrtosiphon pisum</i> (Harris)	AP	Lacotte et al. (2023)
<i>Teucrium polium</i> (L.)	<i>Callosobruchus maculatus</i> (Fab.)	LE	Arab et al. (2008)
<i>Thymus hirtus</i> subsp. <i>algeriensis</i> (Boiss. & Reut.) Murb.	<i>Tribolium castaneum</i> (H) Herbst	AP	Amari et al. (2022)
<i>Thymus algeriensis</i> Boiss. & Reut.	<i>Aphis fabae</i> Scopoli	LE / FL	Kouache et al. (2024)
<i>Bupleurum fruticosum</i> (L)	<i>Spodoptera littoralis</i> (Bois.)	LE/ RO/ ST	Ben El Hadj Ali et al. (2015)
	<i>Sitophilus oryzae</i> (L)	NM	Koutsaviti et al. (2018)
<i>Daucus carota</i> (L.)	<i>Spodoptera littoralis</i> (Bois.)	SE	Ben-Khalifa et al. (2018)
<i>Ferula communis</i> (L.)	<i>Spodoptera littoralis</i> (Bois.)/ <i>Myzus persicae</i> (Sulzer)	NM	Mamoci et al. (2012)
<i>Eryngium tricuspdatum</i> (L.)	<i>Tribolium confusum</i> (J. Du Val)	NM	Majdoub et al. (2024)
<i>Visnaga daucoides</i> Gaertn.	<i>Culex quinquefasciatus</i> Say	NM	Pavela (2008)
<i>Foeniculum vulgare</i> Mill.	<i>Sitophilus granarius</i> (Linnaeus) / <i>Tribolium castaneum</i> (L)	NM	Teke et al. (2021)
	<i>Rhopalosiphum padi</i> (Linnaeus)	LE	Benddine et al. (2023)
	<i>Spodoptera littoralis</i> (Bois.)	SE	Ben-Khalifa et al. (2018)

Species	Pest	Plant part	References
	<i>Acyrtosiphon pisum</i> (Harris)	SE	Lacotte et al. (2023)
	<i>Aphis gossypii</i> Glover	NM	Atanasova et al. (2018)
<i>Thapsia garganica</i> (L.)	<i>Tribolium Castaneum</i> (H) Herbst	SE	Khemis et al. (2023)
	<i>Sitophilus granarius</i> (L)	ST/LE	Chelef et al. (2021)
<i>Smyrniium olusatrum</i> (L.)	<i>Tetranychus urticae</i> Koch	FL	Benelli et al. (2017)
<i>Olea europaea subsp. Europaea</i>	<i>Ephestia kuehniella</i> Zeller	ST	Lahcene et al. (2018)
<i>Asphodelus microcarpus</i> Rchb.	<i>Tribolium castaneum</i> (H) Herbst	AP	Saada et al. (2010)
<i>Urtica dioica</i> (L)	<i>Culex pipiens</i> (Linnaeus)	NM	Toubal et al. (2019)
<i>Myrtus communis</i> (L.)	<i>Callosobruchus maculatus</i> (Fab.), <i>Tribolium confusum</i> (J. Du Val)	LE	Khani et al. (2012)
	<i>Acyrtosiphon pisum</i> (Harris)	AP	Lacotte et al. (2023)
<i>Eucalyptus globulus</i> Labill.	<i>Myzus persicae</i> (Sulzer)	AP	Oulebsir-Mohandkaci et al. (2015)
	<i>Acyrtosiphon pisum</i> (Harris)	LE	Lacotte et al. (2023)
<i>Eucalyptus camaldulensis</i> Dehnh.	<i>Oryzaephilus surinamensis</i> (Linnaeus) / <i>Sitophilus oryzae</i> (L)	LE	Ebadollahi et al. (2020)
	<i>Ectomyelois ceratoniae</i>	LE	Haouel et al. (2010)
	<i>Aphis gossypii</i> Glover	LE	Koorki et al. (2018)
<i>Juniperus oxycedrus</i> (L.)	<i>Sitophylus oryzae/tribolium confusum</i> (J)	NM	Athanassiou et al. (2013)
	<i>Rhopalosiphum padi</i> (Linnaeus) / <i>Sitobion avenae</i> (Fab.) / <i>Fusarium spp</i> / <i>Botrytis cinerea</i> Pers. / <i>Colletotrichum</i> <i>spp.</i> / <i>Rhizoctonia solani</i> Kühn/ <i>Cylindrocarpon pauciseptatum</i> Wollenw.	NM	Semerdjieva et al. (2021)
	<i>Aphis spiraeicola</i> Patch	LE	Amokrane et al. (2024)
<i>Tetraclinis articulata</i> (Vahl) Mast.	<i>Sitophilus granarius/ Tribolium confusum</i> (J. Du Val)	FR	Sadiki et al. (2022)
<i>Laurus nobilis</i> (L.)	<i>Tribolium confusum</i> (J. Du Val)	NM	Isikber et al. (2006)
	<i>Acyrtosiphon pisum</i>	LE	Lacotte et al. (2023)
<i>Pistacia lentiscus</i> (L.)	<i>Lobesia botrana</i> (Denis)	FR/LE	Dasenaki et al. (2022)
	<i>Anopheles maculatus</i> Theobald	FR /LE	Pourya et al. (2018)
	<i>Aphis spiraeicola</i> Patch / <i>Aphis gossypii</i> Glover	NM	Behi et al. (2019)
<i>Convolvulus lineatus</i> (L)	<i>Chaitophorus leucomelas</i> Koch	LE	Zaid et al. (2021)
	<i>Sitophilus oryzae</i> (L)	AP	Allali et al.

Species	Pest	Plant part	References
			(2010)
<i>Bellardia trixago</i> (L.) All	<i>Tribolium castaneum</i> (H) Herbst	AP	Pascual-Villalobos et al. (1998)
<i>Pteridium aquilinum</i> (L.) Kuhn	<i>Chaitophorus populialbae</i>	LE	Zaid et al. (2022)
<i>Cistus creticus</i> (L.)	<i>Fusarium moniliforme</i> / <i>Fusarium culmorum</i> <i>Schltl</i> / <i>Alternaria alternata</i> (Fr.) Keissl / <i>Botrytis cinerea</i> Pers.	AP	Çaşkurlu et al. (2020)
<i>Cistus salviifolius</i> (L.)	<i>G.citri-aurantii</i>	NM	Karim et al. (2017)
	<i>Plodia interpunctella</i> (Hübner)	AP	Bouayad et al. (2013)
<i>Cistus monpeiliensis</i> (L.)	<i>G.citri-aurantii</i>	NM	Karim et al. (2017)
	<i>Plodia interpunctella</i> (Hübner)	AP	Bouayad et al. (2013)
<i>Hypericum perforatum</i> (L.)	<i>Leptinotarsa decemlineata</i>	WP	Daraban et al. (2021)
<i>Hypericum humifusum</i> (L.)	<i>Culex pipiens</i> (Linnaeus)	AP	Rouis et al. (2013)
<i>Raphanus raphanistrum</i> (L.)	<i>Tribolium castaneum</i> (H) Herbst	NM	Jbilou et al. (2008)
<i>Hirschfeldia incana</i> (L.) Lagr.-Foss.	<i>Verticillium dahliae</i> Kleb	LE/ST	Varo et al. (2017)
<i>Verbascum sinuatum</i> (L.)	<i>Aphis gossypii</i> Glov.	WP	Soliman et al. (2005)
<i>Daphne gnidium</i> (L.)	<i>Sitophilus granarius</i> (L)	ST/LE	Chelef et al. (2021)
<i>Rubus ulmifolius</i> Schott	<i>Ceratitidis capitata</i> (Wiedemann)	LE	Ghabbari et al. (2018)
<i>Reseda alba</i> (L.)	<i>Phthorimaea operculella</i> (Zeller)	AP	Mazrou et al. (2021)
<i>Viola odorata</i> (L.)	<i>Acalyptis pistaciae</i> van Nieuwerkerken & Hull	ST/LE/FL	Razavi et al. (2015)
	<i>Ceratitidis capitata</i> (Wiedemann)	LE	Ibrahim et al.2023
<i>Capparis spinosa</i> (L.)	<i>Tribolium confusum</i> (J. Du Val)	FR	Manesh et al. (2022)
	<i>Bemisia tabaci</i> (Gennadius)	SE	Shahrour et al. (2024)
<i>Cytisus villosus</i> Pourr.	<i>Culex pipiens</i> (Linnaeus)	LE	Djidel et al. (2018)
<i>Ceratoniasiliqua</i> (L.)	<i>Fusarium oxysproum</i> f.sp.c / <i>Monilinia fructigena</i> Honey	LE	Yavuz et al. (2018)
<i>Melilotus officinalis</i> (L.) Lam.	<i>Diaphorina citri</i> Kuwayama / <i>Aphis gossypii</i> Glover	NM	Tayyab et al. (2022)
<i>Silene vulgaris</i> (Moench) Garcke	<i>Spodoptera littoralis</i> (Bois.)	ST	Pavela (2011)
<i>Linum bienne</i> (L.)	<i>Rhyzopertha dominica</i> (Fab.)	SE	Moon et al. (2021)
	<i>Myzus persicae</i> (Sulzer)	SE	Haidar et al. (2023)
<i>Verbena officinalis</i> (L.)	<i>Culex pipiens</i> (Linnaeus)	LE	Zoubiri et al.

Species	Pest	Plant part	References
			(2011)(b)
<i>Taxus baccata</i> (L.)	<i>Sitophilus oryzae</i> (L)	LE	Riffi et al. (2019)
<i>Polygonum aviculare</i> (L.)	<i>Tetranychus cinnabarinus</i> (Bois.)	NM	Jin and al. (2012)
<i>Cedrus atlantica</i> (Endl.) Manetti ex Carrière	<i>Dactylopius opuntiae</i> (Cockerell)	LE	El Finti et al. (2022)
<i>Pinus halepensis</i> Mill.	<i>Aphis spiraecola</i> Patch / <i>Fusarium verticillioides</i> (Sacc.) / <i>Fusarium oxysporum</i> Schldl. / <i>Fusarium poae</i> / <i>Fusarium nygamai</i> / <i>Fusarium solani</i> (Mart.) Sacc. / <i>Fusarium redolens</i> Wollenw. / <i>Bipolaris sorokiniana</i> (Sacc.) Shoemaker, / <i>Phoma</i> sp/ <i>Monographella nivalis</i> (Schaffnit) , <i>Stemphylium</i> ,	LE	Derbali et al. (2024)
<i>Papaver rhoeas</i> (L.)	<i>Hylobius abietis</i> (Linnaeus)	LE	Korczyński et al. (2007)
<i>Euphorbia peplus</i> (L.)	<i>Gonipterus platensis</i> (Marelli)	NM	Huerta et al. (2022)
<i>Euphorbia bupleuroides</i> Desf.	<i>Tribolium castaneum</i> (H) Herbst / <i>Trogoderma granarium</i> (Everts)	ST	Brik et al. (2021)
<i>Ricinus communis</i> (L.)	<i>Macrosiphum rosae</i> (Linnaeus)	LE	El Haddad et al. (2021)
	<i>Plutella xylostella</i> (Linnaeus)	LE/RO/SE	Kodjo et al. (2011)
	<i>Tuta absoluta</i> (Meyrick)	NM	Ait Taadaouit et al. (2012)
<i>Mercurialis annua</i> (L.)	<i>Tribolium confusum</i> (J. Du Val)	LE	Ben Nasr et al. (2021)
<i>Aristolochia fontanesii</i> Boiss. & Reut.	<i>Oryzaephilus surinamensis</i> (Linnaeus)	NM	Tail et al. (2017)
<i>Plantago lanceolata</i> (L.)	<i>Spodoptera littoralis</i> (Bois.)	ST	Pavela (2011)
<i>Plantago lagopus</i> (L.)	<i>Rhizoctonia napaeae</i> Westend.	NM	Behiry et al. (2022)
<i>Scrophularia canina</i> (L.)	<i>Culex pipiens</i> (Linnaeus)	AP	Giacinto et al. (2011)
<i>Malva sylvestris</i> (L.)	<i>Colletotrichum lindemuthianum</i> (Sacc. & Magnus)	LE	Pinto et al. (2010)
<i>Althaea officinalis</i> (L.)	<i>Alternaria</i> spp	LE	Maghsoodi et al. (2021)
<i>Sambucus ebulus</i> (L.)	<i>Tribolium confusum</i> (J. Du Val)	LE	Haghighian et al. (2005)
<i>Scabiosa arenaria</i> Forssk.	<i>Culex pipiens</i> (Linnaeus)	WP/FR	Hlila et al. (2017)
<i>Borago officinalis</i> (L.)	<i>Spodoptera littoralis</i> (Bois.)	LE	Hazaa et al. (2020)
	<i>Myzus persicae</i> (Sulzer)	LE	Fawaz et al. (2020)
<i>Nerium oleander</i> (L.)	<i>Chaitophorus leucomelas</i> Koch	LE	Zaid et al. (2021)
<i>Oxalis pes-caprae</i> (L.)	<i>Macrosiphum rosae</i> (L) / <i>Gynaikothrips ficorum</i> (Marchal)	LE/FL	Lachguer et al. (2021)

LE: Leaves, RO: Roots, WP: whole plant, ST: stem, FL: Flowers, AP: Arial partt, SE: Seeds, FR: Fruits, NM: not mentioned




Figure 2. Diagram of the distribution of botanical pesticides by families in the Blidian Atlas




Figure 3. Biological types of the species inventoried




Figure 4. Spectrum of phytogeological repartition of plants inventoried

Table 2. Shannon diversity indices in the different habitats


	F.T	M.T	W.T	F.ME	M.ME	W.ME	F.S	M.S	W.S
Taxa_S	16	40	41	24	55	47	25	55	42
Shannon_H	3.24	4.18	4.20	3.66	4.5	4.34	3.7	4.5	4.23
Veg-zone		3.87			4.17			4.141	

F.T = forests thermo-Mediterranean, M.T = maquis thermo-Mediterranean, W.T = wastlands Thermo-Mediterranean, F.ME = forests meso-Mediterranean, M.ME = maquis meso-Mediterranean, W.ME = wastlands meso-Mediterranean, F.S = forests supra-Mediterranean, M.S = maquis supra-Mediterranean, W.S = wastlands supra-Mediterranean

### Classification

The Ascending Hierarchical Classification (AHC) using the presence-absence Jaccard index identified six groups of species (Fig. 5). The group 1: comprised 52 species, including: *Ajuga iva*, *Althaea officinalis*, *Anacyclus clavatus*, *Aristolochia*

*paucinervis*, *Atractylis cancellata*, *Atractylis gummifera*, *Bellardia trixago*, *Carthamus lanatus*, *Cerantonia siliqua*, *Clinopodium nepeta*, *Clinopodium vulgare*, *Convolvulus lineatus*, *Coleostephus myconis*, *Daucus carota*, *Daphne gnidium*, *Dittrichia viscosa*, *Echinops spinosissimus*, *Erigeron canadensis*, *Eryngium tricuspdatum*, *Euphorbia bupleuroides*, *Eucalyptus camaldulensis*, *Eucalyptus globulus*, *Euphorbia peplus*, *Ferula communis*, *Hypericum humifusum*, *Hypericum perforatum*, *Juniperus oxycedrus*, *Lavandula stoechas*, *Linum bienne*, *Malva sylvestris*, *Mercurialis annua*, *Melilotus officinalis*, *Olea europaea*, *Oxalis pes-caprae*, *Papaver rhoeas*, *Pallenis spinosa*, *Pinus halepensis*, *Pistacia lentiscus*, *Plantago lagopus*, *Plantago lanceolata*, *Reseda alba*, *Rubus ulmifolius*, *Sambucus ebulus*, *Salvia verbenaca*, *Senecio vulgaris*, *Silene vulgaris*, *Sonchus asper*, *Sonchus oleraceus*, *Urtica dioica*, *Verbena officinalis*, *Verbascum sinuatum*, and *Viola odorata*.



**Figure 5.** Classification of the 88 species using the presence absence data and Jaccard similarity index

Group 2 consisted of 9 species: *Borago officinalis*, *Calendula arvensis*, *Capparis spinosa*, *Glebionis coronaria*, *Hirschfeldia incana*, *Mentha rotundifolia*, *Raphanus raphanistrum*, *Ricinus communis*, and *Visnaga daucoides*.

Group 3, the least diverse, contained only two species: *Helichrysum stoechas* and *Myrtus communis*.


Group 4, the second largest, included the following species: *Asphodelus microcarpus*, *Bupleurum fruticosum*, *Cistus creticus*, *Cistus monspeliensis*, *Cistus salviifolius*, *Cytisus villosus*, *Mauranthemum paludosum*, *Origanum floribundum*, *Polygonum aviculare*, *Pteridium aquilinum*, *Satureja montana*, *Scabiosa arenaria*, *Scrophularia canina*, *Smyrniolum olusatrum*, *Teucrium polium*, *Thymus algeriensis*, *Thapsia garganica*, and *Thymus hirtus*.

The fifth and sixth groups each contained three species: *Foeniculum vulgare*, *Nerium oleander*, *Tetraclinis articulata*, and *Cedrus atlantica*, *Laurus nobilis*, and *Taxus baccata*.

### Correspondence analysis (CA)

To explore the relationship between species, habitats, and vegetation zones, we conducted a Correspondence Analysis (CA). Species were coded using the first two letters of the genus and the first two letters of the species. The results revealed that the first two axes (F1 and F2) explained 50.71% of the variance (Fig. 6), facilitating the identification of four distinct groups, each represented by a set of species sharing

similar habitat. The first axis (F1) explained 27.94% of the variance, allowing for the differentiation of group 1 and group 2. Group 1 which was positively correlated with Axis F1, primarily consists of species from the forest habitat. The species in this group include: *Sonchus asper* (Soas), *Viola odorata* (Viod), *Ajuga iva* (Ajiv), *Salvia verbenaca* (Save), *Ceratonia siliqua* (Cesi), *Eucalyptus globulus* (Eugl), *Rubus ulmifolius* (Ruul), *Sonchus oleraceus* (Sool), *Eucalyptus camaldulensis* (Euca), *Senecio vulgaris* (Sevu), *Hypericum perforatum* (Hype), *Hypericum humifusum* (Hyhu), *Pallenis spinosa* (Pasp), *Verbena officinalis* (Veof), *Atractylis gummifera* (Atgu), *Atractylis cancellata* (Atca), *Satureja montana* (Samo), *Taxus baccata* (Taba), *Cedrus atlantica* (Ceat), *Laurus nobilis* (Lano), *Helichrysum stoechas* (Hest), *Myrtus communis* (Myco), *Juniperus oxycedrus* (Juox), *Pinus halepensis* (Piha), *Olea europaea* (Oleu), *Daphne gnidium* (Dagn), and *Pistacia lentiscus* (Pile). Group 2, which was negatively correlated with axis 1 (F1), was primarily composed of species found in the wasteland and maquis habitats across the three vegetation zones: Thermo-Mediterranean, Meso-Mediterranean, and Supra-Mediterranean. This group includes the following species: *Reseda alba* (Real), *Linum bienne* (Libi), *Papaver rhoeas* (Parh), *Nerium oleander* (Neol), *Foeniculum vulgare* (Fovu), *Tetraclinis articulata* (Tear), *Anacyclus clavatus* (Ancl), *Oxalis pes-caprae* (Oxpc), *Malva sylvestris* (Masy), *Verbascum sinuatum* (Vesi), *Silene vulgaris* (Sivu), *Convolvulus lineatus* (Coli), *Euphorbia peplus* (Eupe), *Euphorbia bupleuroides* (Eubu), *Bellardia trixago* (Betr), *Plantago lanceolata* (Plln), *Plantago lagopus* (Pllg), *Urtica dioica* (Urdu), *Daucus carota* (Daca), *Carthamus lanatus* (Cala), *Dittrichia viscosa* (Divi), *Echinops spinosissimus* (Ecsp), *Sambucus ebulus* (Saeb), *Melilotus officinalis* (Meo), *Ferula communis* (Feco), and *Eryngium tricuspidatum* (Ertr).



**Figure 6.** Plot (F1-F2) of the correspondence analysis (CA) conducted on a contingency table of 88 species, three habitats (Forest, Maquis, and Wasteland), and three vegetation zones (Thermo-Mediterranean, Meso-Mediterranean, and Supra-Mediterranean)

The second axis (F2) explained 22.7% of the variance and was positively correlated with Group 3, which consists of the following species: *Visnaga daucooides* (Vida), *Borago officinalis* (Boof), *Glebionis coronaria* (Glco), *Calendula arvensis* (Caar), *Raphanus raphanistrum* (Rara), *Capparis spinosa* (Casp), *Mentha rotundifolia* (Mero), *Hirschfeldia incana* (Hiin), *Ricinus communis* (Rico), and *Coleostephus myconis* (Comy). The species in this group were primarily found in the wastelands of the Thermo-Mediterranean zone. This axis was negatively correlated with Group 4, which was primarily found in the maquis of the Meso-Mediterranean and Supra-Mediterranean zones. The species in this group include: *Thapsia garganica* (Thga), *Althaea officinalis* (Alof), *Cistus creticus* (Cicr), *Cistus monspeliensis* (Cimo), *Cistus salviifolius* (Cisa), *Bupleurum fruticosum* (Bufr), *Mercurialis annua* (Mean), *Lavandula stoechas* (Last), *Clinopodium vulgare* (Clvu), *Clinopodium nepeta* (Clne), *Aristolochia paucinervis* Pomel (Arfo), *Erigeron canadensis* (Erca), *Pteridium aquilinum* (Ptaq), *Asphodelus microcarpus* (Asmi), *Mauranthemum paludosum* (Mapa), *Scabiosa arenaria* (Scar), *Origanum floribundum* (Orfl), *Smyrniolum olusatrum* (Smol), *Cytisus villosus* (Cyvi), *Scrophularia canina* (Scca), *Thymus hirtus* (Thhi), *Teucrium polium* (Tepo), and *Thymus algeriensis* (Thal).

## Discussion

The Blidian Atlas region, like many Mediterranean areas, is renowned for its rich biodiversity, which arises from a combination of complex ecological factors, including vegetation zones: thermo-Mediterranean, meso-Mediterranean, and supra-Mediterranean, as well as topographical and bioclimatic influences. Despite extensive studies on the chemical composition, medicinal properties, and antibacterial effects of essential oils from plants in Algeria and the Blidian Atlas (Mehalaine et al., 2017), research on botanical pesticides in the region remains limited. This study identified a total of 452 plant species from 74 botanical families. In comparison, a recent study by Ouelmouhoub and Benrima (2021) recorded only 104 taxa from 50 botanical families. Additionally, Aouadj et al. (2020) documented the presence of 25 endemic, 31 rare, and 9 protected taxa in the area.

Across Algeria, Zoubiri and Aoumeur (2011) identified 230 plants with pesticide properties and discussed the methodologies for evaluating their practical use. Among the inventoried species, 88 have been identified as botanical pesticides effective against pests and phytopathogens. The most common families included Asteraceae, Lamiaceae, and Apiaceae. Notable species in the study were *Pistacia lentiscus* (L.), *Daucus carota* (L.), *Eryngium tricuspdatum* (L.), *Ferula communis* (L.), *Smyrniolum olusatrum* (L.), *Thapsia garganica* (L.), *Dittrichia viscosa* (Greuter), *Helichrysum stoechas* (L.), *Sonchus oleraceus* (L.), *Cistus salviifolius* (L.), *Cytisus villosus* (Pourr.), *Clinopodium vulgare* (L.), *Lavandula stoechas* (L.), *Eucalyptus camaldulensis* (Dehnh.), *Olea europaea* var. *sylvestris* (Mill.), *Cedrus atlantica* (Manetti ex Endl.), *Pinus halepensis* (Mill.), *Rubus ulmifolius* (Schott), *Scrophularia auriculata* (L.), and *Daphne gnidium* (L.).

The same botanical families were highlighted by Amokrane et al. (2023) and Fragkouli et al. (2023) as having significant pesticide potential. Acheuk et al. (2022) emphasized the active compounds in these families, such as alkaloids, phenols, essential oils, fatty acids, limonoids, and polyketides, and their mechanisms of action, which include toxicity, anti-feeding effects, and growth regulation. According to Fusar et al.

(2024), the biopesticidal properties of these plants play a crucial role in the Euro-Mediterranean region, especially in countries like Tunisia, Spain, and Turkey.

The Lamiaceae family, which accounts for 14% of the inventory, was particularly notable for its essential oils, which possess antimicrobial and insecticidal properties, as highlighted by Salhi et al. (2010), and Adjou and Aoumanou (2013). Despite the large number of species within this family, its insecticidal potential remains largely underutilized.

Most of the species identified as botanical pesticides were hemicryptophytes (31%). According to Houessou et al. (2012) and Poveda et al. (2020), these species are known for their ability to withstand harsh climatic conditions. They are among the first to emerge in early spring and possess bioactive properties that help protect against certain parasites and infections. In contrast, in the semi-arid region of Chlef (Algeria), Amokrane et al. (2023) identified 47 plant species from 21 botanical families, with Lamiaceae, Asteraceae, Apiaceae, and Cupressaceae being the most prevalent. However, therophytes were the most common in this region, with 21 species predominating.

The species recorded in this area reflect a wide range of biogeographical origins, but the Mediterranean component predominates, making up 49% of the floral heritage. This is consistent with the study area's location within the Mediterranean basin and the findings of Ouelmouhoub and Benrima (2021).

Finally, most of the biopesticides recorded in this study were primarily effective against storage pests, such as *Tribolium castaneum*, *Sitophilus granarius*, *Tribolium confusum*, *Callosobruchus maculatus*, and *Sitophilus oryzae*. These pests pose significant challenges to grain preservation and result in substantial economic losses (Pascual-Villalobos et al., 1999; Haouas et al., 2014; Polatoğlu et al., 2017; Rotundo et al., 2019).

## Conclusion

The Blidian Atlas region boasts a rich and diverse plant life, shaped by its unique ecological features, including a variety of bioclimatic and topographical conditions that foster the growth of many rare and endemic species. These wild plants are vital to the local ecosystem and possess significant therapeutic, aromatic, biopesticidal, and commercial potential. Our study identified 88 pesticidal plants from 38 botanical families, highlighting their importance for sustainable agriculture. The ongoing research into isolating and synthesizing the active components of these plants offers promising prospects for developing commercial biopesticides, which can be utilized to combat harmful pests, contributing to more robust and eco-friendly agricultural practices.

**Acknowledgements.** Authors warmly thank Mr Belkhous Arezki from the general conservation of forests in the National Park of Chrea in Blida for helping identify most of the species.

## REFERENCES

- [1] Acheuk, F., Basiouni, S., Shehata, A. A., Dick, K., Hajri, H., Lasram, S., Yilmaz, M., Emekci, M., Tsiamis, G., Spona-Friedl, M., May-Simera, H., Eisenreich, W., Ntougias, S.

- (2022): Status and Prospects of Botanical Biopesticides in Europe and Mediterranean Countries. – *Biomolecules* 12(2): 311. <https://doi.org/10.3390/biom12020311>.
- [2] Adeniyi, S. A., Orjiekwe, C. L., Ehiagbonare, J. E., Arimah, B. D. (2010): Preliminary phytochemical analysis and insecticidal activity of ethanolic extracts of four tropical plants (*Vernonia amygdalina*, *Sida acuta*, *Ocimum gratissimum* and *Telfaria occidentalis*) against beans weevil (*Acanthscelides obtectus*). – *Int. J. Phys. Sci* 5: 753-762.
- [3] Adjou, E., Aoumanou, M. (2013): Efficacité des extraits de plantes dans la lutte contre les moisissures toxigènes isolées de l'arachide en post-récolte au Bénin. – *Journal of Applied Biosciences* 70(1): 5555. doi:10.4314/jab.v70i1.98755.
- [4] Amokrane, D., Mohammedi, A., EL Haouari, M., Bettahar, S., Ababou, A. (2023): Pesticide Plants: Diversity and Use in Crop Protection in a Southern Mediterranean Area: Case Study of Chlef in North-West Algeria. – *Journal of Chemical Health Risks* 13(4): 119-132.
- [5] Aouadj, S. A., Nasrallah, Y., Hasnaoui, O., Khatir, H. (2020): La flore rare, endémique et menacée des monts de Saida (Algérie). – *Agrobiologia* 11(2): 45-57.
- [6] APG IV. (2016): An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. – *Botanical Journal of the Linnean Society* 181(1): 1-20.
- [7] Fragkouli, R., Antonopoulou, M., Asimakis, E., Spyrou, A., Kosma, C., Zotos, A., Tsiamis, G., Patakas, A., Triantafyllidis, V. (2023): Mediterranean Plants as Potential Source of Biopesticides: An Overview of Current Research and Future Trends. – *Metabolites* 13: 967. <https://doi.org/10.3390/metabo13090967>.
- [8] Fusar Poli, E., Fontefrancesco, M. F. (2024): Use of Medicinal Plants in Middle Atlas of Morocco: Potential Health Risks and Indigenous Knowledge in a Berber Community. – *Encyclopedia* 4: 720-734. <https://doi.org/10.3390/encyclopedia402004>.
- [9] Haouas, D., Cioni, P. L., Halima-Kamel, M. B., Flamini, G., Hamouda, M. H. B. (2014): Chemical constituents and toxicity of essential oils from three Asteraceae plants against *Tribolium confusum*. – *Tunisian Journal of Plant Protection* 9(1): 67-82.
- [10] Houessou, L. G., Teka, A., Oumorou, M., Sinsin, B. (2012): Hemicryptophytes plant species as indicator of grassland state in semi-arid region: Case study of W Biosphere Reserve and its surroundings area in Benin (West Africa). – *International Journal of Biological and Chemical Sciences* 6(3): 1271-1280.
- [11] Isman, M. B. (2006): Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. – *Annual Review of Entomology* 51(1): 45-66. doi:10.1146/annurev.ento.51.11010.
- [12] Lengai, G. M. W., Muthomi, J. W., Mbega, E. R. (2019): Phytochemical activity and role of botanical pesticides in pest management for sustainable agricultural crop production. – *Scientific African* 7: e00239.
- [13] Maghsoodi, F., Taheri, P. (2021): Efficacy of *Althaea officinalis* leaf extract in controlling *Alternaria* spp. pathogenic on Citrus. – *European Journal of Plant Pathology* 161: 799-813.
- [14] Meddour, R. (1994): Contribution à l'étude phytosociologique de la portion centro-orientale du parc national de Chréa. – Thèse de Magister, Université Saad Dahleb - Blida, Faculté des Sciences de la Nature et de la Vie.
- [15] Mehalaine, S., Belfadel, O., Menasria, T., Messaili, A. (2017): Chemical composition and antibacterial activity of essential oils of three medicinal plants from Algerian semi-arid climatic zone. – *Phytothérapie*. doi:10.1007/s10298-017-1143-y.
- [16] Mejdoub, K., Mami, I. R., Belabbes, R., Dib, M., DJabou, N., Tabti, B., Benyelles, N. G., Costa, J., Muselli, A. (2020): Chemical Variability of *Atractylis gummifera* Essential Oils at Three Developmental Stages and Investigation of Their Antioxidant, Antifungal and Insecticidal Activities. – *Current Bioactive Compounds* 16(4): 489-497. doi.org/10.2174/1573407215666190126152112.


- [17] Mekideche, S., Brakchi-Ouakour, L., Kadik, L. (2018): Impact des perturbations anthropiques sur la diversité végétale de la subéraie de Chréa, au nord de l'Algérie. – Bois et Forêts des Tropiques 337: 53-66. <https://doi.org/10.19182/bft2018.337.a31633>.
- [18] Neeraj, G. S., Kumar, A., Ram, S., Kumar, V. (2017): Evaluation of nematicidal activity of ethanolic extracts of medicinal plants to *Meloidogyne incognita* (kofoid and white) chitwood under lab conditions. – International Journal of Pure and Applied Bioscience 1: 827-831.
- [19] Ouelmouhoub, S., Benrima, A. (2021): Esquisse botanique dans une subéraie relique de l'Atlas blidéen (région de Chréa, Algérie). – Agrobiologia 11(1): 2430-2443.
- [20] Pascual-Villalobos, M., Robledo, A. (1999): Screening for anti-insect activity in Mediterranean plants. – Industrial Crops and Products 8(3): 183-194. doi:10.1016/s0926-6690(98)00002-8.
- [21] Polatoğlu, K., Karakoç, Ö. C., Demirci, B., Başer, K. H. C. (2017): Chemical composition and insecticidal activity of edible garland (*Chrysanthemum coronarium* L.) essential oil against the granary pest *Sitophilus granarius* L. (Coleoptera). – Journal of Essential Oil Research 30(2): 120-130. doi:10.1080/10412905.2017.140850.
- [22] Poveda, J., Abril-Urías, P., Escobar, C., Li, W. (2020): Biological Control of Plant-Parasitic Nematodes by Filamentous Fungi Inducers of Resistance: Trichoderma, Mycorrhizal and Endophytic Fungi. – Frontiers in Microbiology 11: 992.
- [23] Qin, W., Huang, S., Li, C., Chen, S., Peng, Z. (2010): Biological activity of the essential oil from the leaves of *Piper sarmentosum* Roxb. (Piperaceae) and its chemical constituents on *Brontispa longissima* (Gestro) (coleoptera: hispididae). – Pestic. Biochem. and Physio 96(3): 132-139. doi.org/10.1016/j.pestbp.2009.10.006.
- [24] Quézel, P., Santa, S. (1962-1963): Nouvelle flore de l'Algérie et des régions désertiques méridionales. – Centre National de la Recherche Scientifique (éds), Paris.
- [25] Raunkiaer, C. (1934): The life forms of plants and statistical plant geography. – Oxford: Clarendon Press.
- [26] Rotundo, G., Paventi, G., Barberio, A., De Cristofaro, A., Notardonato, I., Russo, M. V., Germinara, G. S. (2019): Biological activity of *Dittrichia viscosa* (L.) Greuter extracts against adult *Sitophilus granarius* (L.) (Coleoptera, Curculionidae) and identification of active compounds. – Scientific Reports 9: 6429. <https://doi.org/10.1038/s41598-019-42886-4>.
- [27] Salhi, S., Fadli, M., Zidane, L., Douira, A. (2010): Études floristique et ethnobotanique des plantes médicinales de la ville de Kénitra (Maroc). – Lazaroa 31: 133-146. doi:10.5209/rev\_LAZA.2010.v31.9.
- [28] Sande, D., Mullen, J., Wetzstein, M., Houston, J. (2011): Environmental impacts from pesticide use: a case study of soil fumigation in Florida tomato production. – International Journal of Environmental Research and Public Health 12: 4649-4661.
- [29] Sayed, S., Soliman, M. M., Al-Otaibi, S., Hassan, M. M., Elarnnauty, S. A., Abozeid, S. M., El-Shehawi, A. M. (2022): Toxicity, Deterrent and Repellent Activities of Four Essential Oils on *Aphis punicae* (Hemiptera: Aphididae). – Plants 11: 463. <https://doi.org/10.3390/plants11030463>.
- [30] Shabana, Y. M., Abdalla, M. E., Shahin, A. A., El-Sawy, M. M., Draz, I. S., Youssif A. W. (2017): Efficacy of plant extracts in controlling wheat leaf rust disease caused by *Puccinia triticina*. – Egyptian Journal of Basic and Applied Sciences 1: 67-73.
- [31] Shannon, C. E., Weaver, W. (1949): A Mathematical Theory of Communication. – University of Illinois Press.
- [32] Wimalawansa, S. A., Wimalawansa, S. J. (2014): Agrochemical-related environmental pollution: Effects on human health. – Global Journal of Biology, Agriculture and Health Sciences 3: 72-83.
- [33] Zettler, J. L., Cuperus, G. W. (1990): Pesticide resistance in *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Rhyzopertha dominica* (Coleoptera: Bostrichidae) in wheat. – J. Econ. Entomol. 83: 1677-1681.

- [34] Zoubiri, S., Baaliouamer, A. (2011): Larvicidal activity of two Algerian Verbenaceae essential oils against *Culex pipiens*. – *Veterinary Parasitology* 181(2-4): 370-373.
- [35] Zoubiri, S., Baaliouamer, A. (2011): Chemical composition and insecticidal properties of some aromatic herbs essential oils from Algeria. – *Food Chemistry* 129(1): 179-182. <https://doi.org/10.1016/j.foodchem.2011.04.033>.


## APPENDIX


*Appendix of some plant species recorded and their botanical families, some photos showing the sampled areas*

### Lamiaceae



## Asteraceae





### Apiaceae



*Daucus carota* L.



*Ferula communis* L.



*Foeniculum vulgare* Mill.



*Eryngium tricuspdatum* L.



*Bupleurum fruticosum* L.

## Brassicaceae



*Hirschfeldia incana* (L.) Lagr.-Foss.



*Raphanus raphanistrum* L.

### Hypericaceae



*Hypericum humifusum L.*



*Hypericum perforatum L.*

### Asphodelaceae



*Asphodelus microcarpus Rchb.*

### Capparaceae



*Capparis spinosa L.*

### Cistaceae



*Cistus salvifolius L.*

*Cistus monspeliensis L.*



*Cistus creticus L.*


### Plantaginaceae



*Plantago lagopus L.*


*Plantago lanceolata L.*

**Myrtaceae**



*Eucalyptus camaldulensis* Dehnh.

**Oxalidaceae**



*Oxalis pes-caprae* L.

**Orobanchaceae**




*Bellardia trixago* (L.) All

**Fabaceae**



*Cytisus villosus* Pourr.

**Convolvulaceae**



*Convolvulus lineatus* L

**Dennstaedtiaceae**



*Pteridium aquilinum* (L.) Kuhn

**Verbenaceae**



*Verbena officinalis L.*

**Polygonaceae**



*Polygonum aviculare L.*

**Cupressaceae**



*Tetrclinis articulata (Vahl) Mast.*

**Taxaceae**




*Taxus baccata L.*

**Thymelaeaceae**



*Daphne gnidium L.*

### Pinaceae



*Cedrus atlantica (Endl.) Manetti ex Carrière*



*Pinus halepensis Mill.*

### Malvaceae




*Althaea officinalis L.*



*Malva sylvestris L.*

### Resedaceae



*Reseda alba L.*

### Euphorbiaceae



*Mercurialis annua L.*

**Caprifoliaceae**



*Sambucus ebulus L.*

**Papaveraceae**



*Papaver rhoeas L.*

**Lauriaceae**



*Laurus nobilis L.*

**Violaceae**



*Viola odorata L.*

**Aristolochiaceae**



*Aristolochia fontanesii Boiss.*

**Apocynaceae**



*Nerium oleander L.*

### Scrophulariaceae



*Scrophularia canina L.*

#### *Photos showing some sampled areas*



*Meso- mediterranean Maquis*

*Supra- mediterranean Forest*



*Thermo-mediterranean Wasteland*

## APPENDIX 2

### *Bibliographic references in Table 1: Inventoried pesticide plants, used parts and target pests*

- [1] Ait Taadaouit, N., Hsaine, M., Rochdi, A., Nilahyane, A., Bouharroud, R. (2012): Effet des extraits végétaux méthanoliques de certaines plantes marocaines sur *Tuta absoluta* (Lepidoptera, Gelechiidae). – EPPO Bulletin 42(2): 275-280. doi:10.1111/epp.2571.
- [2] Allali, A., Rezouki, S., Louasté, B., Najat, T., Eloutassi, N., Fadli, M. (2020): Investigation of vegetal bio-insecticides for the stored seeds. – International Journal of Botany Studies 5(3): 219-224.
- [3] Alvarez-Castellanos, P., Bishop, C., Pascual-Villalobos, M. (2001): Antifungal activity of the essential oil of flowerheads of garland chrysanthemum (*Chrysanthemum coronarium*) against agricultural pathogens. – Phytochemistry 57: 99-102.
- [4] Amari, R., Guesmi, F., Saidi, I., Bouzenna, H., Khaled, I., Ben Ali, M., Hedfi, A., Alghamdi, A. S., Hfaiedh, N., Allagui, M. S. (2022): Insecticidal and Antimicrobial Potential of the Volatile Oils of the Aerial Parts of *Teucrium ramosissimum* and *Thymus hirtus* subsp. Algeriensis Growing in the South-West of Tunisia. – Journal of Essential Oil-Bearing Plants 25(6): 1185-1207.
- [5] Amokrane, D., Mohammedi, A., Yekhlef, R., Belfennache, D., Daoudi Zerrouki, N., Aly, S. H., Elanany, M. A., Ali, M. A. (2024): Insecticidal Activity, GC/MS Analysis, and in silico Studies of *Juniperus phoenicea* Essential Oil against *Aphis spiraecola*. – Universal Journal of Agricultural Research 12(1): 51-64. DOI: 10.13189/ujar.2024.120106.
- [6] Arab, N. M., Ebadi, R., Hatami, B., Jahromi, K. T. (2008): Insecticidal effects of some plant extracts on *Callosobruchus maculatus* F. under laboratory condition and *Laphygma exigua* H. in greenhouse. – Journal of Science and Technology of Agriculture and Natural Resources 11(42): 221-235.
- [7] Atanasova, D., Ganchev, D., Nevov, N. (2018): Efficacy of some plant essential oils against cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae) under laboratory conditions. – MAYFEB Journal of Agricultural Science 1: 10-16.
- [8] Athanassiou, C. G., Kavallieratos, N. G., Evergetis, E., Katsoula, A.-M., Haroutounian, S. A. (2013): Insecticidal Efficacy of Silica Gel with *Juniperus oxycedrus* ssp. *Oxycedrus* (Pinales: Cupressaceae) Essential Oil against *Sitophilus oryzae* (Coleoptera: Curculionidae) and *Tribolium confusum* (Coleoptera: Tenebrionidae). – Journal of Economic Entomology 106(4): 1902-1910. Doi: 10.1603/ec12474.
- [9] Azeem, M., Zaman, T., Abbasi, A. M., Abid, M., Mozūratis, R., Alwahibi, M. S., Elshikh, M. S. (2022): Pesticidal potential of some wild plant essential oils against grain

- pests *Tribolium castaneum* (Herbst, 1797) and *Aspergillus flavus* (Link, 1809). – Arabian Journal of Chemistry 15(1): 103482.
- [10] Behi, F., Bachrouch, O., Boukhris-Bouhachem, S. (2019): Insecticidal Activities of *Mentha pulegium* L., and *Pistacia lentiscus* L., Essential Oils against Two Citrus Aphids *Aphis spiraecola* Patch and *Aphis gossypii* Glover. – Journal of Essential Oil-Bearing Plants 22(2): 516-525. doi:10.1080/0972060x.2019.1611483.
- [11] Behiry, S. I., Al-Askar, A. A., Soliman, S. A., Alotibi, F. O., Basile, A., Abdelkhalek, A., Elsharkawy, M. M., Salem, M. Z. M., Hafez, E. E., Heflish, A. A. (2022): *Plantago lagopus* extract as a green fungicide induces systemic resistance against Rhizoctonia root rot disease in tomato plants. – Front. Plant Sci 13: 966929. Doi: 10.3389/fpls.2022.96692.
- [12] Belabbes, R., Mami, I. R., Dib, M. E. A., Mejdoub, K., Tabet, B., Costa, J., Muselli, A. (2020): Chemical Composition and Biological Activities of Essential Oils of *Echinops spinosus* and *Carlina vulgaris* Rich in Polyacetylene Compounds. – Current nutrition & Food Science 16(4): 563-570.
- [13] Ben El Hadj Ali, I., Chaouachi, M., Bahri, R., Chaieb, I., Boussaïd, M., Harzallah-Skhiri, F. (2015): Chemical composition and antioxidant, antibacterial, allelopathic and insecticidal activities of essential oil of *Thymus algeriensis* Boiss. et Reut. – Industrial Crops and Products 77: 631-639. doi:10.1016/j.indcrop.2015.09.046.
- [14] Ben Nasr, R., Baudelaire, E. D., Dicko, A., El Ferchichi, O. H. (2021): Phytochemicals, Antioxidant Attributes and Larvicidal Activity of *Mercurialis annua* L. (Euphorbiaceae) Leaf Extracts against *Tribolium confusum* (Du Val) Larvae (Coleoptera; Tenebrionidae). – Biology 10(4): 344.
- [15] Benddine, H., Zaid, R., Babaali, D., Daoudi-Hacini, S. (2023): Biological activity of essential oils of *Myrtus communis* (Myrtaceae, Family) and *Foeniculum vulgare* (Apiaceae, Family) on open fields conditions against corn aphids *Rhopalosiphum maidis* (Fitch, 1856) in western Algeria. – J. Saudi Soc. Agric. Sci 22(2): 78-88.
- [16] Benelli, G., Pavela, R., Canale, A., Nicoletti, M., Petrelli, R., Cappellacci, L., Galassi, R., Maggi, F. (2017): Isofuranodiene and germacrone from *Smyrniololus olusatrum* essential oil as acaricides and oviposition inhibitors against *Tetranychus urticae*: impact of chemical stabilization of isofuranodiene by interaction with silver triflate. – Journal of Pest Science 90: 693-699.
- [17] Ben-Khalifa, N. E., Chaieb, I., Laarif, A., Haouala, R. (2018): Insecticidal activity of six Apiaceae essential oils against *Spodoptera littoralis* Biosduval (Lepidoptera: Noctuidae). – Journal of New Sciences 55(1): 3603-3609.
- [18] Bouayad, N., Rharrabe, K., Ghailani, N. N., Jbilou, R., Castañera, P., Ortego, F. (2013): Insecticidal effects of Moroccan plant extracts on development, energy reserves and enzymatic activities of *Plodia interpunctella*. – Spanish Journal of Agricultural Research 11(1): 189-198.
- [19] Brik, I., Naama, F. (2021): Euphorbia bupleuroides Desf. Latex as biopesticide against the red flour beetle (*Tribolium castaneum* [Herbst, 1797]) and khapra beetle (*Trogoderma granarium* Everts, 1898). – Acta Agriculturae Slovenica 117(2): 1-6.
- [20] Çaşkurlu, A., Karadağ, A., Köse, Y. B., Tosun, F. (2020): In vitro Antifungal Activity of *Cistus creticus* L. against Plant Pathogenic Fungi. – Acta Pharm. Sci. 58(4-2): 413-419. DOI: 10.23893/1307-2080.APS.05822.
- [21] Chelef, M., Hemida, H., Hassani, A., Mazrou, K. (2021): Efficacy of three vegetal powders against *Sitophilus granarius* in stored wheat and the evaluation of their effect on grain parameters. – International Journal of Environmental Studies 78(4): 679-695.
- [22] Daraban, G. M., Badeanu, M., Suteu, D., Asachi, G. (2021): Repellent and insecticide activities of plants extracts from spontaneous flora using conventional and innovative assisted extraction techniques. – Research Journal of Agricultural Science 53(3): 85-92.
- [23] Dasenaki, L., Betsi, P. C., Raptopoulos, D., Konstantopoulou, M. (2022): Insecticidal Effect of *Pistacia lentiscus* (Anacardiaceae) Metabolites against *Lobesia botrana*

- (Lepidoptera: Tortricidae). – Agronomy 12(4): 755.  
<https://doi.org/10.3390/agronomy12040755>.
- [24] Debbabi, H., El Mokni, R., Chaieb, I., Nardoni, S., Maggi, F., Caprioli, G., Hammami, S. (2020): Chemical Composition, Antifungal and Insecticidal Activities of the Essential Oils from Tunisian *Clinopodium nepeta* subsp. *Nepeta* and *Clinopodium nepeta* subsp. *Glandulosum*. – *Molecules* 25(9): 2137. Doi:10.3390/molecules25092137.
- [25] Derbali, F., Hammami, S. T., Algabr, M., Elaieb, M. T., Hamrouni, L. (2024): Chemical composition, Insecticidal and antifungal activities of *Pinus halepensis* mill. and *Acacia cyanophylla* sp. Wood tars. – *Heliyon* 10: e27813.
- [26] Djidel, A., Daghbouche, S., Djazouli, Z. E., Benrima, A. (2020): Evaluation of the insecticidal activity of the crude aqueous extract of the Fabaceae *Cytisus villosus* Pourr. With respect to the dipteran *Culex pipiens* Linnaeus, 1758. – *IOBC/WPRS Bulletin* 151: 119-126.
- [27] Ebadollahi, A., Setzer, W. N. (2020): Analysis of the Essential Oils of *Eucalyptus camaldulensis* Dehnh. and *E. viminalis* Labill. as a Contribution to Fortify Their Insecticidal Application. – *Natural Product Communications* 15(9): 1934578X2094624. Doi:10.1177/1934578x20946248.
- [28] El Finti, A., El Boullani, R., Zahidi, A., El Mousadik, A. (2022): Management of the Prickly Pear Mealy Bug, *Dactylopius opuntiae* Using Bio-Insecticide in Morocco. – *Advances in Entomology* 10: 267-274. <https://doi.org/10.4236/ae.2022.1040>.
- [29] El Haddad, D., Toubal, S., Bouchenak, O., Boumaza, S., Merah, M., Yahiaoui, K., Arab, K. (2022): Insecticidal activity of Algerian castor extracts *Ricinus communis* against the green rose aphid *Macrosiphum rosae*. – *Agrobiologia* 12(1): 2827-2833.
- [30] Fawaz, S. F., Al-Zurfi, S. M., Almulla, A. M. N., Al-Shakry, E. F. H. (2020): Effect of Borage (*Borage officinalis*) Leaves Extract against Borage *officinalis* Aphids in Protected Cultivation. – *Indian Journal of Ecology* 47(12): 275-280.
- [31] Germinara, G. S., Frontera, A. M., De Cristofaro, A., Rotundo, G. (2011): Insecticidal activity of different extracts from *Scrophularia canina* L. against *Culex pipiens molestus* Forskal (Diptera, Culicidae). – *Journal of Environmental Science and Health, Part B*. 46(6): 473-479.
- [32] Ghabbari, M., Guarino, S., Caleca, V., Saiano, F., Sinacori, M., Baser, N., Mediouni-Ben Jemâa, J., Lo Verde, G. (2018): Behavior-modifying and insecticidal effects of plant extracts on adults of *Ceratitis capitata* (Wiedemann) (Diptera Tephritidae). – *Journal of Pest Science* 91(2): 907-917. Doi:10.1007/s10340-018-0952-6.
- [33] Ghalbane, I., Alahyane, H., Aboussaid, H., Chouikh, N., Costa, J., Romane, A., El Messoussi, S. (2022): Chemical Composition and Insecticidal Properties of Moroccan *Lavandula dentata* and *Lavandula stoechas* Essential Oils Against Mediterranean Fruit Fly, *Ceratitis capitata*. – *Neotropical Entomology* 51: 628-636.
- [34] Gonzalez-Coloma, A., Martin-Benito, D., Mohamed, N., Garcia Vallejo, M. C., Soria, A. C. (2006): Antifeedant effects and chemical composition of essential oils from different populations of *Lavandula luisieri* L. – *Biochemical Systematics and Ecology* 34(8): 609-616.
- [35] Gueribis, F., Zermane, N., Khalfi-Habess, O., Siafa, A., Cimmino, A., Boari, A., Evidente, A. (2019): Bioefficacy of compounds from *Dittrichia viscosa* (Asteraceae) as protectant of chickpea seeds against the cowpea seed beetle *Callosobruchus maculatus* (Coleoptera: Chrysomelidae). – *Journal of Plant Diseases and Protection* 126: 437-446. doi:10.1007/s41348-019-00240.
- [36] Haghghian, F., Jalali, J. (2005): Antifeedant, Growth Regulatory and Ovicidal Effect of *Sambucus ebulus* L. Extract on *Tribolium confusum* Duv. – *Caspian J. Env. Sci.* 3(2): 159-162.
- [37] Haidar, H., Akram, A., Ali, A. (2023): Efficacy of different concentrations of flax plant oil *Linum usitatissimum* in controlling green peach aphid *Myzus persicae* (Hemiptera: Aphididae). – *Revis Bionatura* 8(4): 58. <http://dx.doi.org/10.21931/RB/2023.08.04.58>.

- [38] Haouas, D., Cioni, P. L., Halima-Kamel, M. B., Flamini, G., Hamouda, M. H. B. (2014): Chemical constituents and toxicity of essential oils from three Asteraceae plants against *Tribolium confusum*. – Tunisian Journal of Plant Protection 9(1): 67-82.
- [39] Haouel, S., Mediouni-Ben Jemâa, J., Khouja, M. L. (2010): Postharvest control of the date moth *Ectomyelois ceratoniae* using eucalyptus essential oil fumigation. – Tunisian J. Plant Prot 5: 201-212.
- [40] Hazaa, M., Alm-Eldin, M., Ibrahim, A.-E., Elbarky, N., Salama, M., Sayed, R., Sayed, W. (2020): Biosynthesis of Silver Nanoparticles using *Borago officinalis* leaf extract, characterization and larvicidal activity against cotton leaf worm, *Spodoptera littoralis* (Bosid). – International Journal of Tropical Insect Science. doi:10.1007/s42690-020-00187-8.
- [41] Hazafa, A., Jahan, N., Zia, M. A., Rahman, K. U., Sagheer, M., Naeem, M. (2022): Evaluation and optimization of nanosuspensions of *Chrysanthemum coronarium* and *Azadirachta indica* using Response Surface Methodology for pest management. – Chemosphere 292: 133411.
- [42] Hlila, M. B., Lamari, A., Hichri, A. O., Ben Jannet, H., Mastouri, M., Aouni, M., Selmi B. (2017): Larvicidal efficacy of *Scabiosa arenaria* Forssk. (Dipsacaceae) organs extracts against *Culex pipiens* L. – Asian Pac J Trop Dis 7(12): 783-787.
- [43] Huerta, A., Chiffelle, Í., Arias, C., Curkovic, T., Araya, J. E. (2022): Bioefficacy of Euphorbia peplus latex as an antifeedant and insecticide against *Gonipterus platensis* larvae on *Eucalyptus globulus*. – New Zealand Journal of Forestry Science 53: 2. <https://doi.org/10.33494/nzjfs532023x195x>.
- [44] Ibrahim, A. M. A., Soliman, N. A., Alamin, S. M., Mesbah, A. E., Mahmoud A. M. A. (2023): Susceptibility of the different stages of the medfly *Ceratitis capitata* Wiedemann (Diptera: Tephritidae) to the extracts of *Viola odorata* and *Eucalyptus camaldealensis*. – Egyptian Journal of Zoology 80(80): 18-34. DOI: 10.21608/ejz.2022.169125.1089.
- [45] Isikber, A. A., Alma, M. H., Kanat, M., Karci, A. (2006): Fumigant toxicity of essential oils from *Laurus nobilis* and *Rosmarinus officinalis* against all life stages of *Tribolium confusum*. – Phytoparasitica 34(2): 167-177.
- [46] Jbilou, R., Amri, H., Bouayad, N., Ghailani, N., Ennabili, A., Sayah, F. (2008): Insecticidal effects of extracts of seven plant species on larval development,  $\alpha$ -amylase activity and offspring production of *Tribolium castaneum* (Herbst) (Insecta: Coleoptera: Tenebrionidae). – Bioresource Technology 99(5): 959-964. doi:10.1016/j.biortech.2007.03.01.
- [47] Jin, Y. S., Li, Q., Cheng, J., Ma, L., Liu, Y., Wang, Y., Shi, G. (2012): Acaricidal Activities of the Polygonum Aviculare Extracts against *Tetranychus cinnabarinus* and Its Affection to the Enzyme Activities. – Information Tech. And Agricultural Eng AISC. 134: 731-738.
- [48] Karim, H., Boubaker, H., Askarne, L., Cherifi, K., Lakhtar, H., Msanda, F., Boudyach, E. H., Aoumar, A. A. B. (2017): Use of *Cistus aqueous* extracts as botanical fungicides in the control of Citrus sour rot. – Microb. Pathog. 104: 263-267.
- [49] Khani, A., Basavand, F. (2012): Chemical Composition and Insecticidal Activity of Myrtle (*Myrtus communis* L.) Essential Oil against Two Stored-Product Pests. – Journal of Medicinal Plants and By-products 2: 83-89.
- [50] Khemis, E., Mustapha, M. B., Chaieb, I., Ascrizzi, R., Flamini, G., Harrath, A. H., Jannet, H. B., Zardi-Bergaoui, A. (2023): Chemical Composition and Insecticidal Activity against *Tribolium Castaneum* of *Thapsia garganica* L. Seed Essential Oil. – Chemistry and Biodiversity 20(2). <https://doi.org/10.1002/cbdv.202200646>.
- [51] Kodjo, T. A., Gbénonchi, M., Sadate, A., Komi, A., Yaovi, G., Dieudonné, M., Komla, S. (2011): Bio-insecticidal effects of plant extracts and oil emulsions of *Ricinus communis* L. (Malpighiales: Euphorbiaceae) on the diamondback, *Plutella xylostella* L. (Lepidoptera: Plutellidae) under laboratory and semi-field conditions. – J. Appl. Biosci. 43: 2899-2914.

- [52] Koorki, Z., Shahidi-Noghabi, S., Mahdian, M., Pirmaoradi, K. (2018): Chemical Composition and Insecticidal Properties of Several Plant Essential Oils on the Melon Aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae). – Journal of Essential Oil-Bearing Plants 21(2): 420-429. DOI: 10.1080/0972060X.2018.1435308.
- [53] Korczyński, I., Kuźmiński, R. (2007): The effect of oil extracts of selected plant species on feeding of large pine weevil *Hylobius abietis* (L.) beetles on scots pine shoots. – Acta Sci. Pol. Silv. Colendar. Rat. Ind. Lignar 6(2): 5-9.
- [54] Kouache, B., Kaci, Z., Brada, M., Fauconnier, M. L. (2024): Chemical Composition and Insecticidal Activity of *Thymus algeriensis* Boiss and Reut. Essential Oil against *Aphis fabae* Scopoli 1763. – Indian Journal of Agricultural Research 58(1): 164-169.
- [55] Koutsaviti, A., Antonopoulou, V., Vlassi, A., Antonatos, S., Michaelakis, A., Papachristos, D. P., Tzakou, O. (2018): Chemical composition and fumigant activity of essential oils from six plant families against *Sitophilus oryzae* (Col: Curculionidae). – Journal of Pest Science 91: 873-886.
- [56] Lachguer, K., Zakriti, A., Chafik, Z. (2021): Insecticidal effect of some Moroccan plants extracts on *Macrosiphum rosae* and *Gynaikothrips ficorum*. – Journal of Pharmacognosy and Phytochemistry 10(5): 118-124.
- [57] Lacotte, V., Rey, M., Peignier, S., Mercier, P. E., Rahioui, I., Sivignon, C., Razya, L., Benhamoud, S., Livi, S., Da Silva, P. (2023): Bioactivity and chemical composition of forty plant essential oils against the pea aphid *Acyrtosiphon pisum* revealed peppermint oil as a promising biorepellent. – Industrial Crops & Products 197: 116610.
- [58] Lahcene, S., Taibi, F., Mestar, N., Ahmed, S. A., Boumendjel, M., Ouafi, S., Houali, K. (2018): Insecticidal effects of the *Olea europaea* subsp. *Laperrinei* extracts on the flour Pyralid *Ephestia kuehniella*. – Cellular and Molecular Biology 64(11): 6-12.
- [59] Majdoub, S., Chaabane-Banaoué, R., El Mokni, R., Bagané, M., Chaieb, I., Dhaouadi, H., Piras, A., Babba, H., Porcedda, S., Hammam, S. (2024): Chemical Analyses, Antifungal and Insecticidal Properties of the Essential Oils of *Eryngium tricuspdatum* L. (Apiaceae) Growing Natively in Tunisia. – Chemistry Africa 7: 3049-3058.
- [60] Mamoci, E., Cavoski, I., Andres, M. F., Díaz, C. E., Gonzalez-Coloma, A. (2012): Chemical characterization of the aphid antifeedant extracts from *Dittrichia viscosa* and *Ferula communis*. – Biochemical Systematics and Ecology 43: 101-107. doi:10.1016/j.bse.2012.02.012.
- [61] Manesh, T. M., Ravan, S., Khani, A., Sarani, M. (2022): Chemical composition and insecticidal effect of methanol extract of *Capparis spinosa* L. fruits on *Tribolium confusum* Jacquelin du Val, 1863 and *Sitophilus oryzae* (L., 1763) adult. – Acta Agriculturae Slovenica 118(4): 1-6.
- [62] Mazrou, K., Labdelli, F., Bousmaha, F., Chelef, M., Adamou-Djerbaoui, M. (2021): The efficacy of three different plant extracts on some biological features of the potato tuber moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae) by different application methods. – International Journal of Environmental Studies 78(6): 1-12. doi:10.1080/00207233.2021.190224.
- [63] Mejdoub, K., Mami, I. R., Belabbes, R., Dib, M., DJabou, N., Tabti, B., Benyelles, N. G., Costa, J., Muselli, A. (2020): Chemical Variability of *Atractylis gummifera* Essential Oils at Three Developmental Stages and Investigation of Their Antioxidant, Antifungal and Insecticidal Activities. – Current Bioactive Compounds 16(4): 489-497. doi.org/10.2174/1573407215666190126152112.
- [64] Moon, B., Razzaq, M., Haseeb, M., Iftikhar, S., Mushtaq, M., Taimur, M., Shabir, R. (2021): Insecticidal and growth regulatory effect of *Jatropha curcas* and *Linum usitatissimum* extracts along with new chemistry pesticide spinetoram against Sargodha strain of *Rhyzopertha dominica*. – GSC Biological and Pharmaceutical Sciences 17(03): 137-143.
- [65] Mouas, Y., Bouamra, A., Boussad, F., Benrebiha, F. Z., Chaouia, C. (2021): Effet Aphicide De L'extrait Méthanolique D'origanum Floribondum Munby (Lamiaceae) Vis

- Avis Du Puceron des Céréales Sitobion Avenae (aphididae). – *Agrobiologia* 11(2): 2706-2714.
- [66] Navarro-Rocha, J., Andrés, M. F., Díaz, C. E., Burillo, J., González-Coloma, A. (2020): Composition and biocidal properties of essential oil from pre-domesticated Spanish *Satureja Montana* L. – *Ind. Crops Prod.* 145: 111958.
- [67] Oulebsir-Mohandkaci, H., Ait Kaki, S., Doumandji-Mitiche, B. (2015): Essential Oils of two Algerian aromatic plants *Thymus vulgaris* and *Eucalyptus globulus* as Bio-insecticides against aphid *Myzus persicae* (Homoptera: Aphididae). – *Wulfenia Journal* 22(2): 185-197.
- [68] Pavela, R. (2008): Larvicidal effects of various Euro-Asiatic plants against *Culex quinquefasciatus* Say larvae (Diptera: Culicidae). – *Parasitology Research* 102: 555-559.
- [69] Pavela, R. (2011): Screening of Eurasian plants for insecticidal and growth inhibition activity against *Spodoptera littoralis* larvae. – *African Journal of Agricultural Research* 6(12): 2895-2907.
- [70] Pinto, J. M., Souza, E. A., Oliveira, D. F. (2010): Use of plant extracts in the control of common bean anthracnose. – *Crop Protection* 29(8): 838-842.  
doi:10.1016/j.cropro.2010.03.006.
- [71] Pourya, M., Sadeghi, A., Ghobari, H., Taning, C. N. T., Smagghe, G. (2018): Bioactivity of *Pistacia atlantica* desf. Subsp. *Kurdica* (Zohary) Rech. F. and *Pistacia khinjuk* Stocks stocks essential oils against *Callosobruchus maculatus* (Coleoptera: Bruchidae) under laboratory conditions. – *J. Stored Prod. Res* 77: 96-105.
- [72] Razavi, S. H., Mahdian, K. (2015): Evaluation the toxicity of *Viola odorata* extract and Spirotetramat pesticide on the *Agonoscena pistaciae* (Hemiptera: Psyllidea). – *Journal of Entomology and Zoology Studies* 3(5): 110-114.
- [73] Riffi, O., Amechrouq, A., Elhourrin, M., El Idrissi, M., Fliou, J. (2019): Study of the insecticidal activity of *taxus baccata* of the ifrane valley on *Sitophilus oryzae* (L.) (Coleoptera: curculionidae). – *Fresenius Environmental Bulletin* 28(4A): 3197-3202.
- [74] Righi, A. F., Kada, R., Khelil, A., Pujade-Villar, J. (2014): Biological Control against the Cowpea Weevil (*Callosobruchus Chinensis* L., Coleoptera: Bruchidae) Using Essential Oils of Some Medicinal Plants. – *Journal of Plant Protection Research* 54(3): 211-217.  
Doi:10.2478/jppr-2014-0032.
- [75] Rouis, Z., Laamari, A., Abid, N., Elaissi, A., Cioni, P. L., Flamini, G., Aouni, M. (2013): Chemical composition and larvicidal activity of several essential oils from *Hypericum* species from Tunisia. – *Parasitology Research* 112: 699-705.
- [76] Saada, I., Mahdi, K., Boubekka, N., Benzitoun, N., Salhi, O. (2022): Variability of insecticidal activity of *Cupressus sempervirens* L., *Juniperus phoenicea* L., *Mentha rotundifolia* (L.) Huds, and *Asphodelus microcarpus* Salzm. & Viv. Extracts according to solvents and extraction systems. – *Biochemical Systematics and Ecology* 105: 104502.
- [77] Sadiki, F. Z., Bouymajane, A., Sbiti, M., Channaoui, S., Micalizzi, G., Cacciola, F., Dugo, P., Mondello, L., El Idrissi, M. (2022): Chemical profile, antibacterial, antioxidant and insecticidal properties from *Tetraclinis articulata* (Vahl) masters cones. – *Journal of Essential Oil Research of the Essential Oil* 35(5): 383-393.
- [78] Samreen, U., Ibrar, M., Ullah, B., Badshah, L. (2014): Insecticidal activity of *sonchus asper* root and leaf. – *Curr. Opin. Agric.* 3(1): 22-25.
- [79] Semerdjieva, I., Zheljaskov, V. D., Radoukova, T., Dincheva, I., Piperkova, N., Maneva, V., Astatkie, T., Kačániová, M. (2021): Biological Activity of Essential Oils of Four Juniper Species and Their Potential as Biopesticides. – *Molecules* 26(21): 6358.
- [80] Sertkaya, E., Kaya, K., Soylu, S. (2010): Acaricidal activities of the essential oils from several medicinal plants against the carmine spider mite (*Tetranychus cinnabarinus* Boisd.) (Acarina: Tetranychidae). – *Industrial Crops and Products* 31(1): 107-112.
- [81] Shahrour, W. G., Shatnawi, M. A., Al-Alawi, M., Shibli, R. A., Alqudah, T. S., Majdalawi, M. M., Al-Tawaha, A. R., Aljammal, A. (2024): In vitro multiplication,

- antimicrobial, and insecticidal activity of of *Capparis spinosa*. – Not Bot Horti Agrobi 52(1): 13609.
- [82] Shamer, R., El-Shazly, M. M., Khider, A. M., Kordy, A. M. (2020): Chemical Composition of Five Botanical Powders and Their Insecticidal Activity against the Rice Weevil, *Sitophilus oryzae* on Wheat Crop. – Egypt. Acad. J. Biolog. Sci. (G. Microbiolog) 12(1): 125-136.
- [83] Shehzad, M. Z., Riaz, M. A., Raza, A. B. M., Majeed, M. Z. (2021): Biocidal Potential of Indigenous Flora of Soon Valley (Khush-ab, Pakistan) against *Helicoverpa armigera* Hübner and *Spodoptera litura* Fabricius (Lepidoptera: Noctuidae). – In Proceedings of the 1st International Electronic Conference on Entomology. Doi:10.3390/IECE-10368.
- [84] Soliman, M. M. M., Hassanein, A. A., Abou-yousef, H. M. (2005): Efficiency of Various Wild Plant Extracts against the Cotton Aphid, *Aphis gossypii* Glov. (Aphididae: Homoptera). – Acta Phytopathologica et Entomologica Hungarica 40(1-2): 185-196.
- [85] Tail, G., Cherif, H. S., Boughendja, A., Tair, A. (2017): Insecticide activity of *Aristolochia longa* aqueous extract against *Oryzaephilus surinamensis* adults. – 6e COMAPPI, Conférence sur les Moyens Alternatifs de Protection pour une Production Intégrée, Lille, France, pp. 432-437 ref. 21.
- [86] Tayyab, M. B., Majeed, M. Z., Riaz, M. A., Aqueel, M. A., Ouedraogo, S. N., Luqman, M., Ahmed, K. S., Tanvir, M. (2022): Insecticidal potential of indigenous flora of soon valley against Asian citrus Psyllid *Diaphorina citri* kuwayama and cotton aphid *Aphis gossypii* glover. – Sarhad Journal of Agriculture 38(1): 26-39. DOI. <https://dx.doi.org/10.17582/journal.sja/2022/38.1.26.39>.
- [87] Teke, M. A., Mutlu, Ç. (2021): Insecticidal and behavioral effects of some plant essential oils against *Sitophilus granarius* L. and *Tribolium castaneum* (Herbst). – Journal of Plant Diseases and Protection 128: 109-119.
- [88] Toubal, S., Elhaddad, D. J., Bouchenak, O., Yahiaoui, K., Sadaoui, N., Arab, K. (2019): L'importance des extraits d'*Urtica dioica* L. dans la lutte contre *Culex pipiens* (Linné). – Algerian Journal of Environmental Science and Technology 5(1): 868-872.
- [89] Varo, A., Mulero-Aparicio, A., Adem, M., Roca, L. F., Raya-Ortega, M. C., López-Escudero, F. J., Trapero, A. (2017): Screening water extracts and essential oils from Mediterranean plants against *Verticillium dahliae* in olive. – Crop Protection 92: 168-175. doi:10.1016/j.cropro.2016.10.018.
- [90] Wagner, L. S., Sequin, C. J., Foti, N., Campos-Soldini, M. P. (2021): Insecticidal, fungicidal, phytotoxic activity and chemical composition of *Lavandula dentata* essential oil. – Biocatalysis and Agricultural Biotechnology 35: 102092. doi:10.1016/j.bcab.2021.102092.
- [91] Wimalawansa, S. A., Wimalawansa, S. J. (2014): Agrochemical-related environmental pollution: Effects on human health. – Global Journal of Biology, Agriculture and Health Sciences 3: 72-83.
- [92] Yakhlef, G., Hambaba, L., Pinto, D. C. G. A., Silva, A. M. S. (2020): Chemical composition and insecticidal, repellent and antifungal activities of essential oil of *Mentha rotundifolia* (L.) from Algeria. – Industrial Crops and Products 158: 112988.
- [93] Yavuz, A., Onaran, A., Yanar, Y. (2018): Antifungal Activity of Leaf Extract of *Ceratonia siliqua* L. Plant against Plant Pathogenic Fungi. – International Journal of Agricultural and Natural Sciences 1(1): 36-38.
- [94] Zaid, R., Canela-Garayoa, R., Ortega-Chacon, N. M., Mouhouche, F. (2022): Phytochemical analyses and toxicity of *Nerium oleander* (Apocynaceae) leaf extracts against *Chaitophorus leucomelas* Koch, 1854 (Homoptera: Aphididae). – Journal of Saudi Society of Agricultural Science 21(5): 310-317.
- [95] Zaid, R., Mouhouche, F., Canela-Garayoa, R., Benddine, H., Ortega Chacón, N. M. (2022): Chemical composition and insecticidal activity of two eagle fern (*Pteridium aquilinum* (L.) Kuhn) extracts on the poplar aphid *Chaitophorus populialbae* (Hemiptera-Aphididae). – J. Cent. Eur. Agric. 23(4): 773-781.

**APPENDIX 3**

*Table of existence absence of plant species in the three habitats and the three vegetation stages*

Species	T	M	S	FT	FM	FS	MT	MM	MS	WT	WM	WS
Ancl	1	1	0	0	0	0	1	1	0	1	1	0
Atgu	1	1	1	1	1	1	0	0	0	1	1	1
Atca	1	1	1	1	1	1	0	0	0	1	1	1
Caar	1	0	0	0	0	0	0	0	0	1	0	0
Cala	1	1	1	0	0	0	1	1	1	1	1	1
Comy	1	1	0	0	0	0	0	0	0	1	1	0
Divi	1	1	1	0	0	0	1	1	1	1	1	1
Ecsp	1	1	1	0	0	0	1	1	1	1	1	1
Erca	0	1	1	0	0	0	0	1	1	0	1	1
Glco	1	0	0	0	0	0	0	0	0	1	0	0
Hest	0	1	0	0	1	0	0	1	0	0	0	0
Mapa	0	1	1	0	0	0	0	1	1	0	0	0
Pasp	1	1	1	1	1	1	1	1	1	1	1	1
Soas	0	1	1	0	1	1	0	1	1	0	1	1
Sool	1	1	1	1	1	1	1	1	1	1	1	1
Sevu	1	1	1	1	1	1	1	1	1	1	1	1
Ajiv	0	1	1	0	1	1	0	1	1	0	1	1
Clne	0	1	1	0	0	0	0	1	1	0	1	1
Clvu	0	1	1	0	0	0	0	1	1	0	1	1
Last	0	1	1	0	0	0	0	1	1	0	1	1
Lade	0	0	1	0	0	0	0	0	1	0	0	1
Mero	1	0	0	0	0	0	0	0	0	1	0	0
Saver	0	1	1	0	1	1	0	1	1	0	1	1
Orfl	0	1	1	0	0	0	0	1	1	0	0	0
Samo	0	0	1	0	0	1	0	0	1	0	0	0
Tepo	0	0	1	0	0	0	0	0	1	0	0	0
Thhi	0	0	1	0	0	0	0	0	1	0	0	0
Thal	0	0	1	0	0	0	0	0	1	0	0	0
Bufr	1	1	1	0	0	0	1	1	1	0	0	0
Daca	1	1	1	0	0	0	1	1	1	1	1	1
Feco	0	1	1	0	0	0	0	0	0	0	1	1
Ertr	0	1	1	0	0	0	0	0	0	0	1	1
Vida	1	0	0	0	0	0	0	0	0	1	0	0
Fovu	1	0	0	0	0	0	1	0	0	0	0	0
Thga	0	0	1	0	0	0	0	0	0	0	0	1
Smol	0	0	1	0	0	0	0	0	1	0	0	0
Oleu	1	1	1	1	1	1	1	1	1	0	0	0
Asmi	0	1	1	0	0	0	0	1	1	0	0	0
Urdu	1	1	1	0	0	0	1	1	1	1	1	1
Myco	0	1	0	0	1	0	0	1	0	0	0	0

Species	T	M	S	FT	FM	FS	MT	MM	MS	WT	WM	WS
Eugl	1	1	1	1	1	1	1	1	1	1	1	1
Euca	1	1	1	1	1	1	1	1	1	1	1	1
Juox	0	1	1	0	1	1	0	1	1	0	0	0
Tear	1	0	0	0	0	0	1	0	0	0	0	0
Lano	0	1	1	0	1	1	0	0	0	0	0	0
Pile	1	1	1	1	1	1	1	1	1	0	0	0
Coli	1	1	1	0	0	0	1	1	1	1	1	1
Betr	1	1	1	0	0	0	1	1	1	1	1	1
ptaq	0	1	1	0	0	0	0	1	1	0	0	0
Cicr	1	1	1	0	0	0	1	1	1	0	0	0
Cisa	1	1	1	0	0	0	1	1	1	0	0	0
Cimo	1	1	1	0	0	0	1	1	1	0	0	0
Hype	1	1	1	1	1	1	1	1	1	1	1	1
Hyhu	1	1	1	1	1	1	1	1	1	1	1	1
Rara	1	0	0	0	0	0	0	0	0	1	0	0
Hiin	1	0	0	0	0	0	1	0	0	1	0	0
Vesi	1	1	0	0	0	0	1	1	0	1	1	0
Dagn	1	1	1	1	1	1	1	1	1	0	0	0
Ruul	1	1	1	1	1	1	1	1	1	1	1	1
Real	1	1	0	0	0	0	1	1	0	1	1	0
Viod	0	1	1	0	1	1	0	1	1	0	1	1
Casp	1	0	0	0	0	0	0	0	0	1	0	0
Cyvi	0	1	1	0	0	0	0	1	1	0	0	0
Cesi	1	1	1	1	1	1	1	1	1	1	1	1
Meo	0	1	1	0	0	0	0	0	0	0	1	1
Sivu	1	1	1	0	0	0	1	1	1	1	1	1
Libi	1	1	1	0	0	0	0	0	0	1	1	1
Veof	1	1	1	1	1	1	1	1	1	1	1	1
Taba	0	0	1	0	0	1	0	0	0	0	0	0
Poav	0	0	1	0	0	0	0	0	1	0	0	1
Ceat	0	0	1	0	0	1	0	0	0	0	0	0
Piha	1	1	1	1	1	1	1	1	1	0	0	0
Parh	1	1	1	0	0	0	0	0	0	1	1	1
Eupe	1	1	1	0	0	0	1	1	1	1	1	1
Eubu	1	1	1	0	0	0	1	1	1	1	1	1
Rico	1	0	0	0	0	0	1	0	0	1	0	0
Mean	0	1	1	0	0	0	0	1	1	0	1	1
Arfo	0	1	1	0	0	0	0	1	1	0	1	1
Pllg	1	1	1	0	0	0	1	1	1	1	1	1
Plln	1	1	1	0	0	0	1	1	1	1	1	1
Scca	0	0	1	0	0	0	0	0	1	0	0	0
Masy	1	1	0	0	0	0	1	1	0	1	1	0
Alof	0	1	0	0	0	0	0	1	0	0	1	0
Saeb	0	1	0	0	0	0	0	0	0	0	1	0

Species	T	M	S	FT	FM	FS	MT	MM	MS	WT	WM	WS
Scar	0	1	1	0	0	0	0	1	1	0	0	0
Boof	1	0	0	0	0	0	0	0	0	1	0	0
Neol	1	0	0	0	0	0	1	0	0	0	0	0
Oxpc	1	1	0	0	0	0	1	1	0	1	1	0

T: Thermo-medeterranean  
 M: Meso-medeterranean  
 S: Supra-medeterranean  
 FT: Forest Thermo-medeterranean  
 FM: Forest Meso-medeterranean  
 FS: Forest supra-medeterranean  
 MT: Maquis Thermo-medeterranean  
 MM: Maquis Meso-medeterranean  
 MS: Maquis supra-medeterranean  
 WT: wasteland Thermo-medeterranean  
 WM: wasteland Meso-medeterranean  
 WS: wasteland supra-medeterranean

Taxa_S	16	40	41	24	55	47	25	55	42
Individuals	16	40	41	24	55	47	25	55	42
Dominance_D	0	0	0	0	0	0	0	0	0
Simpson_1-D	1	1	1	1	1	1	1	1	1
Shannon_H	3.241	4.176	4.201	3.657	4.498	4.34	3.699	4.498	4.226
Evenness_e^H/S	1.598	1.628	1.629	1.615	1.634	1.631	1.616	1.634	1.629
Brillouin	1.917	2.758	2.781	2.283	3.06	2.911	2.32	3.06	2.804
Menhinick	4	6.325	6.403	4.899	7.416	6.856	5	7.416	6.481
Margalef	5.41	10.57	10.77	7.237	13.48	11.95	7.456	13.48	10.97
Equitability_J	1.169	1.132	1.131	1.151	1.123	1.127	1.149	1.123	1.131
Fisher_alpha	0	0	0	0	0	0	0	0	0
Berger-Parker	0.0625	0.025	0.02439	0.04167	0.01818	0.02128	0.04	0.01818	0.02381
Chao-1	128.5	800.5	841	288.5	1513	1105	313	1513	882.5
iChao-1	128.5	800.5	841	288.5	1513	1105	313	1513	882.5
ACE	128.5	800.5	841	288.5	1513	1105	313	1513	882.5

**This paper has also an electronic APPENDIX 4.**