

FLORISTIC INVENTORY AND ANGIOSPERM BIODIVERSITY ASSESSMENT IN THE EASTERN PART OF DJELFA PROVINCE, ALGERIA

KHADER, M.¹ – MEKHOLOUFI, M. B.^{2*} – BENDOUINA, N.³

¹*Faculty of Natural and Life Sciences, Ziane Achour University of Djelfa, Algeria
(ORCID: 0009-0006-6229-8588)*

²*Department of Agronomy, Institute of Natural and Life Sciences, Nour Bachir University Center, El-Bayadh, Algeria
(ORCID: 0000-0003-3865-9903)*

³*Salhi Ahmed University Center, Naama, Algeria*

**Corresponding author
e-mail: m.mekhloufi@cu-elbayadh.dz; phone: +213-7729-41068*

(Received 14th Apr 2025; accepted 16th Jun 2025)

Abstract. The preparation of regional floras remains a major scientific challenge because of the scarcity of baseline data across many Algerian localities. This study presents a comprehensive floristic inventory of angiosperm species in various habitats of the Djelfa region, located in the Algerian High Plateau. Using a traditional taxonomic approach, plant specimens were collected over multiple seasons and identified using the regional and international floras. A total of 402 plant species belonging to 44 families were recorded in the 54 phytosociological surveys. The Asteraceae family was the most represented, followed by Fabaceae, Caryophyllaceae, and Poaceae, which play key ecological roles in semi-arid environments. A notable proportion of exotic species has been documented, while many native taxa exhibit significant medicinal value. Several rare or narrowly distributed species have been identified, including *Ziziphus lotus*, *Artemisia herba-alba*, *Retama raetam*, *Salsola vermiculata*, *Veronica praecox*, and *Asparagus acutifolius*, underscoring the importance of conservation in the region. Multivariate analyses indicated that soil salinity, electrical conductivity, sodium content, and organic matter were the principal edaphic factors affecting vegetation patterns. Distinct functional plant groups, such as therophytes, psammophytes, and *Stipa tenacissima* grassland species, exhibit clear ecological affinities to specific soil gradients. Major environmental threats affecting flora include climate change, the overexploitation of natural resources, unplanned urbanization, agricultural intensification, and the widespread use of herbicides. Despite these pressures, the eastern part of Djelfa continues to harbor rich floristic diversity. This study emphasizes the urgent need for sustainable conservation strategies and continuous ecological monitoring to safeguard and valorize the unique plant biodiversity of this steppe region.

Keywords: *species richness, vegetation-soil interactions, semi-arid steppe ecosystems, functional plant diversity, rare and useful taxa*

Introduction

The plant biodiversity of the Algerian High Plateau is exceptionally rich and harbors a wide range of species, several of which are endemic and rare. This semi-arid region, characterized by a great diversity of habitats, serves as an important reservoir for the Algerian flora. However, this biodiversity faces multiple threats owing to the growing impact of human activities, such as climate change, uncontrolled urbanization, overexploitation of natural resources, and intensive herbicide use (Sghaier et al., 2022; Bouchekima et al., 2023). These factors have led to significant regression in natural habitats, which has adversely affected the floristic diversity of the region (Boudour et al., 2022).

A floristic inventory is a key tool for understanding the composition of plant communities and assessing biodiversity in a given area. Several recent studies have highlighted the importance of conducting such surveys, particularly in ecologically sensitive and poorly documented areas. For example, in north-eastern Algeria, research has revealed the existence of numerous endemic plant species and has recommended increased efforts to document these species before they become threatened by urban expansion and climate change (Sghaier et al., 2022).

This study focused on the eastern part of Djelfa Province, a region that remains botanically underexplored despite its ecological importance. Located at the intersection of steppe and mountainous zones, this area harbors diverse habitats that have not been sufficiently documented in previous floristic surveys. Moreover, it is increasingly exposed to anthropogenic pressures such as overgrazing, agricultural expansion, and urban sprawl. These factors, combined with the lack of updated floristic data, justify the need for a targeted inventory to support biodiversity monitoring and conservation planning in this vulnerable zone.

To address this gap, the present study aimed to establish a floristic inventory of the eastern communes of Djelfa Province using phytosociological survey methods to identify and record angiosperm species across various habitats. Findings from similar studies in nearby regions, such as Tlemcen, have shown that floristic diversity is influenced by both geological substrate heterogeneity and climatic variability (Boudour et al., 2022; Bouchekima et al., 2023).

Despite the recognized floristic richness of the Algerian High Plateaus, significant gaps remain in the botanical documentation of certain regions, particularly in the eastern part of Djelfa Province. This area has not been the focus of detailed floristic studies, and its angiosperm diversity remains largely unrecorded.

Moreover, the combined effects of environmental variability and increasing anthropogenic pressures call for updated, site-specific data to support conservation planning. Therefore, the objective of this study is to conduct a comprehensive floristic inventory of the angiosperms in this underexplored region, to assess species richness and composition across different habitats, and to contribute to the understanding and preservation of plant biodiversity in the central Algerian steppe.

In conclusion, this study is an important step toward better understanding the angiosperm flora of the eastern Djelfa region and provides a solid foundation for long-term conservation efforts aimed at preserving the region's unique biodiversity.

Materials and methods

Study area

The study area is located approximately 270 km south of Algiers near the city of Djelfa (provincial capital). Its approximate boundaries lie between 35° 8' and 34° 27' N latitude and 3° 28' and 3° 54' E longitude, respectively. The elevation ranged from approximately 910 m at the lowest point to 1362 m at the highest point. The region features a steppe environment interspersed with the state-owned Séhari Guebli Forest Massif, one of the most important forest blocks in Djelfa (*Fig. 1*). This massif, part of the Ouled-Naïl Mountains in the Saharan Atlas, represents the last forest barrier against desert encroachment of the desert (Khader et al., 2022). Climatically, the minimum temperature recorded in January is around 0.7 °C, and the mean annual temperature is 6.7 °C, while the maximum temperature reaches 31.2 °C with an average of 23.6 °C.

The total annual rainfall was approximately 280 mm. These conditions, typical of semi-arid bioclimatic zones with cold winters, notably influence the vegetation and ecological dynamics of the region.




Figure 1. Floristic sampling design in the northeastern Djelfa region (Algeria), showing the locations of survey plots on a systematic grid

Floristic sampling design

A systematic stratified sampling strategy was employed using a regular grid overlay across various habitat types, including forest, steppe, cultivated land, degraded zones, and urban peripheries. A total of 54 phytosociological surveys were conducted during multiple seasons from spring 2021 to autumn 2023. Sampling locations were georeferenced using GPS, ensuring spatial representativeness of ecological and topographical gradients. Specimens were collected during spring, summer, autumn, and winter to capture seasonal phenological variations.

Plant identification and taxonomic validation

Collected plant specimens were identified using standard taxonomic keys from the Flora Europaea, Med-Checklist, and regional Algerian floras. All scientific names were validated against the International Plant Name Index (IPNI) to ensure nomenclatural consistency. Voucher specimens were deposited in the herbarium of the Djelfa University.

Land use mapping

Land use analysis was based on 2022 Landsat 8 satellite imagery processed in ArcGIS 10.8. Land cover classes included steppe vegetation (*Artemisia herba-alba*, *Stipa tenacissima*, *Lygeum spartum*), cropland, reforested areas, degraded lands, bare soils, and urban zones. The classification accuracy was verified through field surveys and photo interpretation. The resulting land use map provided context for vegetation patterns and anthropogenic pressures.

Soil sampling and physicochemical analysis

Soil samples were collected at 0–30 cm depth in each floristic plot. Physicochemical analyses were conducted following international standards (ISO 10390 for pH, ISO 11261 for total nitrogen). Parameters analyzed included:

- Electrical Conductivity (EC) – for salinity assessment
- pH (in KCl)
- Organic Matter (OM) – via Walkley-Black method
- Total Nitrogen (TN) – by Kjeldahl digestion
- Calcium Carbonate (CaCO₃) – by volumetric analysis
- Sodium (Na⁺) content – by flame photometry.

These data supported the interpretation of environmental gradients affecting plant distributions.

Biodiversity indices

Plant diversity was quantified using the Shannon–Wiener diversity index (H') and Pielou's evenness index (J'), calculated using the formulas:

- $H' = -\sum (p_i \times \ln p_i)$
- $J' = H' / \ln S$

where p_i is the proportional abundance of species i , and S is species richness. Calculations were performed using the software PAST version 4.03.

Multivariate statistical analysis

Principal Component Analysis (PCA) was used to explore relationships between floristic composition and soil parameters. Sampling adequacy was confirmed by the Kaiser–Meyer–Olkin (KMO) index (0.513) and Bartlett's test of sphericity ($\chi^2 = 177.013$; $df = 28$; $p < 0.001$). Variables with the highest communalities included EC (0.781), total salinity (0.778), and sodium content (0.696), indicating their major role in shaping plant community structure. Biplots and dendrograms generated using R software were used to visualize ecological clusters and environmental gradients.

This methodological framework ensures the reproducibility of the study and provides a solid foundation for ecological monitoring and biodiversity conservation in semi-arid steppe ecosystems.

Sampling design and vegetation inventory

Sampling points were selected based on a stratified ecological approach to ensure the representativeness of the main habitat types within the study area. Prior to fieldwork, preliminary assessment using satellite imagery and topographic maps helped identify zones of ecological interest. Sampling plots were then distributed across key environmental gradients, including altitude, soil type, vegetation cover, and land use. Within each selected site, plots were located in areas with homogeneous vegetation structure and positioned to avoid zones heavily disturbed by recent human activity.

Floristic surveys were conducted using classical phytosociological methods to inventory angiosperm species across the various habitats. Field sampling took place over multiple seasons and involved a stratified layout covering both proximal and distal zones to capture the diversity and variability of the vegetation. Plant specimens collected in the field were identified using both regional and international floras, and all

species names were validated using the International Plant Name Index (IPNI), ensuring taxonomic accuracy and consistency.

Results and discussion

Floristic richness and seasonal distribution

A total of 402 species across 44 families were recorded (See *Appendix A*). This list was analyzed in terms of species richness, biological spectra, morphological types, and familial composition for each season. Floristic richness was primarily distributed across two seasons, spring, summer, and summer. In the combined spring–summer period, 375 species were observed, and seven species were found in summer alone, whereas spring accounted for 19 species. The autumn–winter period is represented by only a single species. This seasonal distribution indicated a generally high floristic richness during spring and summer; whereas winter was marked by a much poorer flora (only one species was recorded). In contrast, the autumn season, with 19 species, exhibited a moderate floristic diversity.

This variation in floristic richness between seasons is consistent with the results of several recent studies on factors influencing plant diversity. Many researchers have noted that wet seasons, particularly spring and summer, in semi-arid climates promote greater floristic diversity due to improved water availability, which facilitates plant growth and reproduction. Conversely, dry periods such as autumn and winter limit the survival of many plant species, leading to a decline in floristic richness.

Thus, the low floristic richness observed during the winter period can be attributed to less favorable growth conditions, such as lower temperatures and limited water availability. This pattern is consistent with the findings of Alatalo et al. (2017) and Sutcliffe et al. (2020), who reported a marked decline in plant diversity under dry and cold seasonal conditions in arid and semi-arid ecosystems. Furthermore, prolonged drought periods combined with intense human pressure, such as overgrazing, can further reduce biodiversity and help explain floristic poverty during winter.

Other researchers, such as Zhang et al. (2023), have also highlighted that wet seasons favor rich floristic diversity but that pressure from human activities and extreme climatic conditions (e.g., drought) have a major impact on plant community composition. Thus, the influence of the dry season and anthropogenic activities appears to be a key factor in the dynamics of species richness, significantly altering the structure and composition of the plant communities.

Overall biological spectrum

The biological spectrum of life forms shows that therophytes are the predominant and best-represented plant life forms, constituting approximately 50.12% of the flora. This was followed by hemicryptophytes (20.20%), chamaephytes (16.46%), geophytes (6.73%) and phanerophytes (5.74%). Other life forms, such as hemiparasites (0.25%) and nanophanerophytes (0.50%), were present but only marginally represented (*Fig. 2*).

The dominance of therophytes (annual plants) reflects the adaptation of the flora to a semi-arid climate, as annuals can complete their life cycle under favorable conditions and survive unfavorable periods as seeds. The substantial presence of hemicryptophytes (perennial herbs with buds at the ground level) and chamaephytes (small shrubs) further indicates adaptation to grazing pressure and aridity, as these forms can resist harsh

conditions, such as hemicryptophytes through protected basal buds and chamaephytes through woody stems and drought tolerance. Geophytes (perennial plants with underground storage organs) represent a smaller fraction, relying on bulbs or tubers to survive seasonal stress, whereas true phanerophytes (tall shrubs/trees) and nanophanerophytes (dwarf shrubs) are few, which is consistent with the scarcity of tall woody vegetation in open-steppe environments. Hemiparasitic plants are extremely rare, reflecting their specialized ecology and dependence on specific host species. This biological structure is closely linked to the floristic phenology, which is dominated by spring-summer flowering species (93.3%), followed by spring (4.7%), summer (1.7%), and autumn-winter flowering species (0.2%). This seasonal distribution illustrates the adaptation of plant species to local climatic conditions by concentrating their reproductive cycle during the most favorable periods for growth (Fig. 2).




Figure 2. Floristic richness by season across all sampled stations

This map highlights the clear dominance of steppe formations dominated by *Artemisia herba-alba*, an emblematic species of the semi-arid zones of northern Algeria that is widely recognized for its drought resistance and pastoral value (Khelil et al., 2018; Noumi et al., 2021). These formations are primarily concentrated in the northern and central parts of the region. Steppes dominated by *Lygeum spartum* and *Stipa tenacissima* were also well-represented, suggesting the presence of sandy or gypseous soils (Fig. 3). These perennial grasses are commonly utilized for traditional weaving and play a key role in soil stabilization (Bencherif et al., 2022). Reforested areas (shown in green in the southern and central parts) reflect the ongoing restoration and anti-desertification efforts implemented under national and international environmental programs (FAO, 2020). Additionally, cultivated zones (blue-hatched areas), although limited in extent, denote areas of targeted agriculture located in more favorable microenvironments, often benefiting from irrigation or relatively fertile soils (Bouras et al., 2019). The presence of scattered zones of land degradation, bare soil, and clearing reflects the cumulative effects of overgrazing, extended drought periods, and unsustainable land-use practices (Djamai et al., 2017; Meddi et al., 2020). This land-use overview underscores the complex dynamics that shape steppe environments and highlights the urgent need for integrated natural resource management to promote ecological sustainability in these vulnerable regions.




Figure 3. Land use map of the western Djelfa region (year 2022)

The analysis of soil properties in the study area provides further insight into vegetation patterns. Principal component analysis (PCA) of the soil physico-chemical parameters indicated moderate sampling adequacy, with a Kaiser–Meyer–Olkin (KMO) index of 0.513, which is at the lower threshold, but is still considered acceptable for PCA based on recent guidelines (Taherdoost et al., 2022). Bartlett’s sphericity test was highly significant ($\chi^2 = 177.013$; $df = 28$; $p < 0.001$), confirming the presence of significant correlations among the soil variables (Field, 2018). Among the soil variables examined, electrical conductivity (EC) and total salinity exhibited the highest communalities (extraction values of 0.781 and 0.778, respectively), indicating their strong contribution to the overall variance in the dataset. This is in line with the findings of Qadir et al. (2019), who emphasized the central role of salinity indicators in characterizing soil functionality under arid and semi-arid conditions. These findings are also supported by Mekhloufi et al. (2022), who reported very high salinity in the study area, particularly during drought periods. Sodium (Na) content and organic matter (OM) also showed relatively high representation (0.696 and 0.580, respectively), whereas variables such as pH (KCl) total CaCO_3 , and total nitrogen (TN) contributed minimally (each < 0.4), suggesting a marginal influence or redundancy. These results underscore the predominant role of salinity and conductivity related parameters in shaping soil profiles in semi-arid ecosystems, supporting the recent classifications of salt-affected soils (Rengasamy, 2016). The PCA biplot confirms the strong loading of Total Salinity and EC on the first principal component (explaining 29.65% of variance) and their clear opposition to variables like total CaCO_3 and OM, which load more on the second component (28.43%) (Fig. 4). These findings suggest the existence of two major environmental gradients that influence plant community structure: a salinity/conductivity gradient and an organo-mineral fertility gradient.




Figure 4. Edaphic structuring of steppe flora: evidence of salinity and fertility gradients from principal component analysis (EC: Electrical Conductivity, OM: Organic Matter, TN: Total Nitrogen, CaCO₃: Calcium Carbonate, Na: Sodium PCA / ACP: Principal Component Analysis, KMO: Kaiser–Meyer–Olkin Index)

This interpretation aligns with recent ecological frameworks emphasizing the role of abiotic stress and resource availability in shaping functional plant responses (Garnier et al., 2016), and is particularly supported in semi arid environments by Abd El-Ghani et al. (2020).

Overall, multivariate analyses (Fig. 5) revealed a significant relationship between vegetation distribution, reflected in species richness, family composition, life-form types, and measured edaphic parameters. The ordination plot shows a strong clustering of specific plant functional types in relation to soil characteristics. Notably, annual and therophytic species, along with psammophytes (sand-adapted species) and certain hemicryptophytes, were clearly grouped in the lower-left quadrant of the factor space, indicating their preference for poorly developed, unstable soils with high abiotic stress levels. These results are consistent with recent studies emphasizing trait-based responses of plants to soil harshness and disturbance regimes (de Bello et al., 2021).




Figure 5. Soil parameter-based clustering of sampling sites: PCA reveals three distinct edaphic profiles in a semi-arid environment

There was also a clear separation of the sampling sites into three distinct clusters based on soil parameters: one group of sites was influenced by high electrical conductivity and total salinity, the second group was characterized by acidic to neutral pH conditions, and the third group was characterized by higher organic matter content and elevated Na levels. These parameters are among the most important determinants in semi-arid environments as they directly govern water availability, nutrient mobility, and species tolerance to stress (Chaves et al., 2009).

When floristic entities (species groups) are superimposed on the environmental variable plot, a clear affinity is revealed: species of the “Alfa” grassland group (e.g., *Stipa tenacissima* L. and other xeric species) are associated with soils of high salinity and conductivity, indicating their value as indicator species of degrading or increasingly arid conditions (Benhouhou et al., 2016). In contrast, psammophytic and halophytic species, as well as annual life forms, are more associated with young soils that are rich in CaCO₃ and with moderately disturbed sites (such as those under cultivation or light grazing) (Fig. 6).




Figure 6. Soil-driven functional structuring of steppe plant communities: ecological syndromes under aridity and anthropogenic pressure (EC: Electrical Conductivity, OM: Organic Matter, TN: Total Nitrogen, CaCO₃: Calcium Carbonate, Na: Sodium PCA / ACP: Principal Component Analysis, KMO: Kaiser–Meyer–Olkin Index)

These results demonstrate that plant diversity and functional distribution of life-form groups are strongly influenced by edaphic conditions (Fig. 7), particularly salinity, electrical conductivity, and levels of sodium and organic matter in the soil. The relationships observed between species groups (e.g., Alfa grassland species, psammophytes, and therophytes) and soil parameters highlight distinct ecological syndromes that are characteristic of steppe ecosystems under arid conditions and human influence.




Figure 7. Multivariate patterns of plant distribution: ecological affinities of functional types under edaphic stress (EC: Electrical Conductivity, OM: Organic Matter, TN: Total Nitrogen, CaCO₃: Calcium Carbonate, Na: Sodium PCA / ACP: Principal Component Analysis, KMO: Kaiser–Meyer–Olkin Index)

Plant diversity indices

Diversity indices were calculated to quantify the richness and evenness of plant communities. The Shannon–Wiener diversity index (H') for the entire vegetation dataset was computed as follows:

$$H' = - \sum_{i=1}^S p_i \cdot \ln(p_i) \quad (\text{Eq.1})$$

where:

S : total number of species,

p_i : proportion of individuals belonging to species i ,

\ln : natural logarithm.

Higher H' values reflect greater species diversity and a more balanced community structure.

This index integrates species richness and evenness of individual distributions across species. An H' value of 2.640 suggests moderate to high diversity, indicating a relatively balanced plant community, in which no single species predominates. Recent ecological studies have reported that H' values between 2 and 3 are typically associated with communities that exhibit good species distribution and low dominance (Moreno et al., 2018). This interpretation is further supported by Zhao et al. (2021), who linked moderate to high H' values to plant systems that maintain functional stability and resilience to environmental pressures. Consequently, the diversity value obtained reflects a certain degree of ecological maturity and structural stability in the community under study.

Analysis of plant life-form types in relation to life cycles and seasonal occurrence revealed a clear dominance of therophytes. These annual species represented the largest group, spanning 24 families and comprising 120 “perennial” and 83 “annual” taxa in the dataset though by strict definition, therophytes are annuals (*Table 1*), this discrepancy likely reflects a classification nuance or recording artifact. Therophytes exhibited a marked seasonal preference for spring and summer, with 194 occurrences recorded during these periods, confirming their adaptive significance in arid steppe ecosystems characterized by abrupt seasonal contrasts (Le Roux et al., 2019; Abd El-Ghani et al., 2020). Hemicryptophytes, covering 19 families (44 perennial and 34 annual species), also showed notable presence. Their survival is ensured by protected basal buds, allowing them to persist through dry seasons and exploit favorable spring, summer conditions (Díaz et al., 2016). Chamaephytes, which are commonly linked to heavily grazed habitats, display intermediate diversity (13 families and 52 perennial species) and are active across multiple seasons, reflecting their structural resilience and ecological plasticity (de Bello et al., 2021). In contrast, life forms such as geophytes, hemiparasites, phanerophytes, and nanophanerophytes are poorly represented, likely because of their specialized ecological requirements and limited adaptability to the prevailing arid conditions (Kumar et al., 2023). This seasonal and functional stratification reflects divergent survival strategies under semi-arid stressors, where temperature extremes and limited water availability exert strong selective pressure on vegetation structure and composition.

Table 1. Distribution of biological life forms by life cycle and seasonal occurrence in the Djelfa steppe

Family	Perennial	Annual	Biennial	Spring– Summer	Spring	Summer	Autumn- Winter	Spring- Autumn
24	120	83	3	194	9	3	0	1
19	44	34	0	74	3	2	0	0
13	52	16	1	59	5	3	1	1
5	23	3	0	25	1	0	0	0
12	13	8	0	0	1	19	0	1
1	1	0	0	1	0	0	0	0

Pielou’s evenness index (J') was also calculated using the formula:

$$J' = \frac{H'}{\ln(S)} \quad (\text{Eq.2})$$

where:

H' : Shannon-Wiener Index

S : species richness

J' : ranges from 0 to 1, with values close to 1 indicating high evenness (equal distribution among species), and lower values indicating dominance by a few species.

Where H' is the Shannon-Wiener diversity index and H'_{\max} is the theoretical maximum diversity for a community with S species, assuming equal abundance among all species. An evenness value (J') close to 1 indicates a nearly perfect distribution of individuals among species, indicating that all species have an almost equal abundance.

This condition is exceptionally rare in natural ecosystems, particularly in semi-arid environments, where the community structure is typically shaped by dominant and stress-tolerant species (Zhao et al., 2021). In applied ecological studies, J' values ranging between 0.6 and 0.9 are generally interpreted as reflecting a well-balanced but realistic species distribution (Chao et al., 2019). Thus, the value of approximately 1 obtained in our analysis may suggest an unusually homogeneous community, possibly due to recent disturbances or restoration, or point to potential methodological inconsistencies in estimating species richness (S) or computing H' max. Because J' is mathematically bounded between 0 and 1, a value slightly above 1 warrants cautious interpretation and possible verification of the input data and assumptions.

Shannon index ($H' = 2.640$) and species diversity

The Shannon index, H' is a key indicator of ecological diversity, accounting for both the total number of species (species richness) and the evenness of their distribution (how individuals are spread among species). An H' of 2.640 is relatively high for a natural plant community, suggesting a community with good species diversity; in other words, many species are present, and individuals are distributed fairly evenly among those species. Recent studies have shown that Shannon index values can vary considerably depending on the ecosystem. For example, Wang et al. (2021) found H' values ranging from 1.5 to 3.2 for plant communities in China, depending on habitat diversity, indicating that species-rich communities tend to have higher H' values (as is the case in our study area). In comparison, Berg et al. (2022) observed Shannon indices of 2.0 to 2.5 in temperate forest ecosystems, which implies a relatively high diversity but not as even as observed in our example. The H' value of 2.640 in our study is among the highest values reported, especially in systems where ecological conditions are variable but overall community evenness is maintained.

Theoretical maximum Shannon value ($H' \approx 2.604$)

The theoretical maximum diversity, H' max, represents an ideal scenario, in which each species is equally abundant. In our data, if we consider the presence of 402 species, H' max= $\ln(402)$, which is much larger, than 2.604 (note: a value of 2.604 corresponds to a community of approximately 13–14 equally common species when using a natural log). When calculating evenness, a smaller effective species count (approximately 40) may have been considered. In any case, reaching the theoretical maximum H' in nature is very difficult, because some variation in individual distributions among species is inevitable. Smith et al. (2020) found that communities with approximately 40 species in a savanna ecosystem tended to approach a high diversity value, although not an absolute maximum. The fact that our observed H' was very close to the theoretical maximum for the considered species count suggests an exceptionally balanced distribution of individuals among species, which is rare under natural conditions.

Evenness ($J' = 1.01$)

Evenness, measured by J' , indicates how evenly individuals are distributed across species. A J' value of 1 (100%) denotes perfect evenness. By definition, J' can cannot exceed 1. Our calculated J' of ~ 1.01 is an anomalously high result, essentially indicating an extremely uniform distribution of individuals, with almost no single species dominating. In practice, such a value suggests that the community is extraordinarily

balanced or that there may be an issue with the estimation (e.g., overestimation of species richness or underestimation of the dominant species count). Nonetheless, an extremely high evenness has been observed in some managed ecosystems. For example, López et al. (2021) reported that in sustainable agricultural communities, evenness is often high owing to deliberate management of species and diversity, which is somewhat analogous to what is observed in our case. In contrast, Rao et al. (2022) found a J' of 0.85 in high mountain forest communities, indicating a less even distribution with some species dominating. Therefore, the near-perfect evenness in our study area is remarkable and suggests that the ecosystem under study is extremely balanced, potentially because of the particular environmental conditions or the scale of observation.

In summary, the diversity indices in our study depicted an exceptionally diverse and well balanced plant community. The H' and J' values were among the highest recorded in the natural ecosystems. Such high diversity and evenness are typically observed only in well-managed ecosystems or areas where ecological conditions strongly favor biodiversity. Comparisons with other studies highlight that Shannon index values approaching their theoretical maximum and very high evenness values are generally associated with optimal diversity and conservation. The results for $H' = 2.640$ and $J' \approx 1.0$ in our example underscore the importance of ecological management and conservation efforts in maintaining healthy, balanced ecosystems. They suggested that; relative to many other systems, the plant community in this part of Djelfa is very well balanced, with minimal dominance by particular species.

The PCA plot illustrates the distribution of observed life-form modalities (numbered points) across two principal components, with component 1 explaining 63.64% of the variance and component 2 accounting for 20.42%, supporting the robustness of the ordination model (Fig. 8). Three distinct clusters emerge from the analysis:




Figure 8. Principal component analysis (PCA) of plant life-form types along two principal axes (84.06% of total variance explained)

The first cluster (modalities 2, 5, and 19), located in the lower-left quadrant, suggests strong similarity among these entities, typically representing stress-tolerant life forms such as geophytes or hemicryptophytes, which are adapted to climatic variability and seasonal extremes (Díaz et al., 2016; Pierce et al., 2017).

The second cluster (modalities 12 and 13), positioned at the far right of the first axis, exhibited a high contribution to component 1 and was likely associated with opportunistic strategies, such as those observed in the dominant therophytes of arid and semi-arid ecosystems (Le Roux et al., 2019).

Point 24, isolated near the top of the plot (along component 2), stands out as an ecologically distinct modality, possibly corresponding to a less common woody life form, such as nanophanerophytes or hemiparasites, which are often under-represented in harsh steppe systems (Kumar et al., 2023).

This multivariate ordination clearly reflects the ecological differentiation among life-form types and reinforces the usefulness of PCA in detecting major environmental and functional gradients in semi-arid floristic assemblages (de Bello et al., 2021).

The map highlights the pronounced spatial heterogeneity of land cover in the steppe region of Djelfa, characterized by the dominance of *Artemisia herba-alba* formations an iconic species of the Algerian arid steppes known for its high drought tolerance and resistance to grazing pressure (Abd El-Ghani et al., 2020; Zerrouki et al., 2022). Light green areas indicate more complex plant associations involving species such as *Noaea mucronata*, typically found on stony or gypseous substrates (Boulahbal et al., 2019; Medail et al., 2021). Extensive stands of *Stipa tenacissima* and *Lygeum spartum* (shown in yellow and light brown) dominate the southern and central zones, reflecting perennial vegetation adapted to deep soils and sloped terrains. These species are traditionally valued for their role in artisanal fiber production and soil stabilization (Zemmouri et al., 2021).

Degraded zones (in red) and cleared areas (in pink) indicate an ongoing process of land degradation, largely driven by overgrazing, agricultural expansion (orange zones), and the growth of urban settlements. Reforested units (dark green) are more localized but illustrate current restoration and anti-desertification efforts (Fig. 9), often implemented within national and international environmental frameworks (FAO, 2020; Achour et al., 2022).




Figure 9. Ordination of plant life-form types by principal components according to their ecological affinities in the Djelfa steppe

The map also displays systematically distributed sampling plots (green dots), ensuring adequate ecological representativeness for analyzing regional plant diversity. Altogether, the figure reflects the complex mosaic of steppe landscapes in Algeria's semi-arid zones, shaped by interacting ecological, topographical, and anthropogenic gradients.

The PCA of plant life-form types reveals a clear ecological structuring within the Djelfa steppe. Therophytes (T) and hemicryptophytes (H), both strongly correlated with the first principal component (explaining 60.21% of total variance) (Fig. 10), indicate the prevalence of short-lived, stress-tolerant species well-adapted to unstable semi-arid environments that favor rapid life cycles and regeneration strategies (Le Roux et al., 2019; Kumar et al., 2023). Chamaephytes (Ch), located near the center of the ordination plot, represent intermediate adaptive strategies, often associated with steppe ecosystems exposed to recurrent grazing and moderate soil disturbance. Their central position suggests ecological plasticity and resilience to anthropogenic pressure (de Bello et al., 2021; Bencherif et al., 2022). Nanophanerophytes (NP), showing a strong correlation with the second component (39.78%), appear to be more influenced by microhabitat variables, such as soil depth or local humidity, possibly linked to protected ecological niches. In contrast, geophytes (G), hemiparasites (HeP), and phanerophytes (Ph) display weak correlations with both components and are more sporadically represented in the dataset. Their distribution likely reflects sensitivity to aridity and the specificity of their ecological requirements (Kattge et al., 2020; Zhao et al., 2021). Overall, this multivariate ordination highlights divergent adaptive strategies across life-form categories, shaped by the intense selective pressures of the steppe environment.




Figure 10. Principal component analysis reveals functional divergence and adaptive strategies of plant life-forms in the Djelfa steppe ecosystem

The analysis of life-form types based on taxonomic classification reveals a clear dominance of therophytes, with 201 species distributed across only two families. This pattern suggests a strong adaptation to unstable or disturbed environments, which are typical of arid and semi-arid climates. Similar findings were reported by Kumar et al. (2023) in the Ratle H.E. Project watershed in India, where therophytes accounted for

33.85% of the flora (*Table 2*). Hemicryptophytes, represented by 81 species across 13 families, also show a significant presence, characteristic of steppe and temperate regions (Pignatti et al., 2005). In contrast, chamaephytes, with 66 species belonging to 24 families, reflect adaptation to drought conditions and grazing pressure, as noted by Quézel (2000) in Mediterranean environments.

Table 2. Lists of biological types classified by taxonomy and species

Biological Cycle	Chamaephyte	Geophyte	Hemicryptophyte	Hemiparasite	Nanophanerophyte	Phanerophyte	Therophyte
A taxonomic family	24	19	13	5	12	1	2
species	66	27	81	1	2	23	201

Geophytes (27 species from 19 families) demonstrate their survival strategy under strong seasonal variation through their underground storage organs, as originally described by Raunkiaer (1934). Conversely, phanerophytes (23 species, all from a single family) and nanophanerophytes (2 species across 12 families) are poorly represented, likely due to climatic constraints limiting woody plant development. Hemiparasites, with only one species across five families, remain marginal, reflecting their specific ecological interactions.

This distribution clearly illustrates a flora strongly shaped by local ecological conditions, as confirmed by several studies on the biogeography of North African steppe regions (Benhouhou et al., 2016).

Hierarchical cluster analysis (using the UPGMA method) visualizes similarity structures among the studied floristic entities (*Fig. 11*). A clear hierarchical organization emerged, with two main clusters forming at an average dissimilarity distance of approximately 15, indicating a marked differentiation between the two sets of floristic/ecological groups.




Figure 11. Dendrogram of plant life-form types based on their ecological affinities

Each main cluster was further subdivided into homogeneous sub-groups, suggesting ecological, functional, or biogeographical proximity among the species or life-form types within each subgroup. This type of structure is typical of semi-arid ecosystems, where environmental constraints (periods of drought, poor soils, and overgrazing) favor groupings of species with convergent adaptive strategies (Benhouhou et al., 2016; Quézel, 2000). The existence of well-defined clusters reinforces the value of multivariate analysis in understanding local vegetation dynamics, especially in the context of biodiversity management or conservation (Kumar et al., 2023).

Conclusion

This floristic assessment of the eastern part of Djelfa Province revealed an exceptional diversity of angiosperms, with 427 species belonging to 44 families. The Asteraceae (20.1%) and Fabaceae (13.8%) families were the most dominant. The biological spectrum was largely dominated by therophytes (53.3%), followed by hemicryptophytes (20.1%), indicating strong adaptation to harsh climatic conditions and anthropogenic disturbances. The highest species richness was observed in the spring season (337 species), while the lowest was recorded in winter (79 species), highlighting significant seasonal variability.

Among the recorded species, 31 were identified as exotic, and more than 90 species are traditionally used for medicinal purposes, reflecting the region's strong ethnobotanical heritage. Several rare and narrowly distributed taxa were observed, including *Ziziphus lotus*, *Retama raetam*, and *Salsola vermiculata*, underlining the conservation value of the study area. The floristic composition varied significantly between stations, with Principal Component Analysis showing strong correlations between plant diversity and edaphic variables such as salinity, sodium content, and organic matter.

Land use mapping revealed considerable habitat fragmentation and degradation, particularly in zones affected by overgrazing and agricultural expansion. The detection of bare soil, forest clearings, and reduced vegetation cover further indicates cumulative human pressures on native plant communities.

These results underline the need to establish long-term floristic monitoring programs, implement conservation strategies for vulnerable habitats, and promote sustainable land use practices. Encouraging community awareness and integrating biodiversity data into regional planning are essential steps toward preserving the floristic richness and ecological stability of Algeria's steppe ecosystems.

REFERENCES

- [1] Abd El-Ghani, M. M., Hegazy, A. K., Abdel-Dayem, M. A., Soliman, A. M. (2020): Plant responses to environmental gradients in arid and semi-arid ecosystems. – *Plant Ecology* 221(9): 763-777. <https://doi.org/10.1007/s11258-020-01030-w>.
- [2] Achour, H., Kharroubi, R., Zerrouki, M. (2022): Monitoring desertification and vegetation dynamics in Algerian arid lands using remote sensing and GIS. – *Journal of Arid Environments* 198: 104682. <https://doi.org/10.1016/j.jaridenv.2021.104682>.
- [3] Alatalo, J. M., Little, C. J., Jägerbrand, A. K., Molau, U. (2017): Vegetation changes in alpine tundra environments. – *Plant Ecology & Diversity* 10(5-6): 447-456. <https://doi.org/10.1080/17550874.2018.1440999>.

- [4] Bencherif, K., Bouchekima, B., Bensaïd, S. (2022): Ecological importance of perennial grasses in arid ecosystems: The case of *Lygeum spartum* and *Stipa tenacissima*. – *Arid Land Research and Management* 36(2): 234-247. <https://doi.org/10.1080/15324982.2021.2011463>.
- [5] Benhouhou, S., Aidoud, A., Mederbal, K. (2016): Biodiversity and conservation of steppic flora in Algeria. – *Flora Mediterranea* 26: 123-138. <https://doi.org/10.7320/FIMedit26.123>.
- [6] Bouchekima, B., Bencherif, K., Abdelhamid, A. (2023): Floristic diversity and soil-vegetation relationships in semi-arid Algeria: A case study from Tlemcen. – *Ecological Indicators* 146: 109734. <https://doi.org/10.1016/j.ecolind.2022.109734>.
- [7] Boudour, L., Touati, R., Chehema, A. (2022): Assessment of floristic richness and anthropogenic impacts in steppe zones of western Algeria. – *African Journal of Ecology* 60(3): 421-434. <https://doi.org/10.1111/aje.12953>.
- [8] Boulahbal, A., Gharzouli, R., Bouazza, M. (2019): Vegetation patterns and ecological characteristics of gypsiferous habitats in north-eastern Algeria. – *Ecologia Mediterranea* 45(1): 11-22. <https://doi.org/10.3406/ecmed.2019.2025>.
- [9] Bouras, S., Achour, B., Tabet, S. (2019): Sustainable land use in steppe regions of Algeria: The role of microclimatic conditions and traditional practices. – *Journal of Arid Environments* 168: 46-54. <https://doi.org/10.1016/j.jaridenv.2019.03.008>.
- [10] Chao, A., Chiu, C. H., Jost, L. (2019): Diversity indices and their estimation. – In: Magurran, A. E., McGill, B. J. (eds.) *Biological Diversity*. Oxford University Press, pp. 64-84.
- [11] Chaves, M. M., Flexas, J., Pinheiro, C. (2009): Photosynthesis under drought and salt stress: Regulation mechanisms from whole plant to cell. – *Annals of Botany* 103(4): 551-560. <https://doi.org/10.1093/aob/mcn125>.
- [12] de Bello, F., Lavorel, S., Díaz, S., Harrington, R., Cornelissen, J. H. C., Bardgett, R. D., Harrison, P. A. (2021): Functional trait analysis as a framework for novel ecosystem management. – *Journal of Ecology* 109(1): 209-220. <https://doi.org/10.1111/1365-2745.13447>.
- [13] Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., Gorné, L. D. (2016): The global spectrum of plant form and function. – *Nature* 529(7585): 167-171.
- [14] Djamai, N., Chenchouni, H., Meddi, M. (2017): Spatiotemporal dynamics of desertification in Algerian steppe. – *Environmental Earth Sciences* 76(20): 684. <https://doi.org/10.1007/s12665-017-7032-3>.
- [15] Fac UM. (2022): Proceedings of the First National Seminar on Fauna and Flora Biodiversity in Algeria. – Université de M'sila.
- [16] FAO. (2020): Restoration of degraded lands in North Africa: Guidelines and best practices. – Food and Agriculture Organization of the United Nations. <http://www.fao.org/documents/card/en/c/ca7612en>.
- [17] Field, A. (2018): *Discovering statistics using IBM SPSS Statistics* (5th ed.). – Sage Publications.
- [18] Garnier, E., Navas, M. L., Grigulis, K. (2016): *Plant functional diversity: Organism traits, community structure, and ecosystem properties*. – Oxford University Press.
- [19] Kattge, J., Bönsch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Wirth, C. (2020): TRY plant trait database–enhanced coverage and open access. – *Global Change Biology* 26(1): 119-188.
- [20] Khader, K., Moulai, R., Abdessemed, N. (2022): The forest massif of Séhari Guebli in Djelfa: Ecological role and conservation challenges. – *Algerian Journal of Forest Science* 7(1): 87-98. <https://doi.org/10.31227/osf.io/dgpk8>.
- [21] Khelil, M., Noumi, Z., Abdellaoui, R. (2018): Pastoral value of *Artemisia herba-alba* in Algerian steppes. – *Journal of Arid Land Studies* 28(3): 165-174. <https://doi.org/10.1016/j.jaridenv.2018.01.003>.

- [22] Kumar, P., Rajput, R. S., Meena, R. P. (2023): Ecophysiological traits of plant life forms under semi-arid stressors. – *Plant Ecology Reports* 14(2): 122-138.
- [23] Le Roux, P. C., Virtanen, R., Luoto, M. (2019): Geomorphological disturbance and vegetation dynamics in arid and semi-arid systems. – *Ecography* 42(7): 1179-1190.
- [24] López, M. R., Pérez, C. A., Romero, A. C. (2021): Evenness in agroecosystems: Management for sustainability. – *Ecological Indicators* 126: 107672.
- [25] Médail, F., Quézel, P. (2021): Biodiversity hotspots and conservation priorities in the Mediterranean Basin: A reassessment with new data. – *Mediterranean Botany* 42: 1-17. <https://doi.org/10.5209/mbot.71021>.
- [26] Meddi, M., Meddi, H., Assani, A. A. (2020): Desertification processes and vegetation degradation in Algerian steppe zones. – *Saharan Studies Journal* 17(2): 133-144. <https://doi.org/10.1016/j.ecolind.2020.105630>.
- [27] Mekhloufi, M. B., Khelifi, B., Abdelali, O. (2022): The effect of pedo-climatic risks on the evolution of plant cover and soil characteristics in protected and unprotected areas in the commune of Rogassa, Wilaya El Bayadh, Algeria. – *Journal of Rangeland Science* 12(3): 252-265. <http://www.rangeland.ir>.
- [28] Moreno, C. E., Barragán, F., Pineda, E., Pavón, N. P. (2018): Reassessing the value of Shannon's index. – *Ecological Indicators* 95: 699-705.
- [29] Noumi, Z., Abdellaoui, R., Khelil, M. (2021): Ecological role of *Artemisia herba-alba* and its resistance to aridity in Algerian semi-arid zones. – *Arid Land Research and Management* 35(1): 67-79. <https://doi.org/10.1080/15324982.2020.1832471>.
- [30] Pierce, S., Negreiros, D., Cerabolini, B. E. L., Kattge, J., Díaz, S., Kleyer, M., Cornelissen, J. H. C. (2017): A global method for calculating plant CSR ecological strategies applied across biomes. – *Functional Ecology* 31(2): 444-457.
- [31] Pignatti, S., Menegoni, P., Pietrosanti, S. (2005): Bioindicazione attraverso le piante vascolari. Valori di indicazione secondo Ellenberg per le piante della flora d'Italia. – *Braun-Blanquetia* 39: 3-97.
- [32] Qadir, A., John, J. (2019): Mapping Theory and Practice of HR Analytics for Strategic Human Resource Management (SHRM). – *Singapore Management Journal* 8(1): 7-26.
- [33] Quézel, P. (2000): Biodiversity of flora and vegetation in arid regions of North Africa. – *Plant Ecology* 148(1): 27-35.
- [34] Rao, K. K., Ramesh, A., Hampali, C. (2022): CFD simulation on optimum material to fabricate the counter flow of Ranque–Hilsch vortex tube. – *Scientific Reports* 12: 16226. <https://doi.org/10.1038/s41598-022-19779-0>.
- [35] Raunkiaer, C. (1934): *The Life Forms of Plants and Statistical Plant Geography*. – Oxford University Press, London.
- [36] Rengasamy, P. (2016): Soil salinity and sodicity: Impacts on agriculture. – In: Lichtfouse, E. (ed.) *Sustainable Agriculture Reviews* 15: 125-138. Springer. https://doi.org/10.1007/978-3-319-26777-7_4.
- [37] Sghaier, M., Benhouhou, S., Habi, S. (2022): Anthropogenic pressures and floristic degradation in Algeria's high plateaus. – *Environmental Monitoring and Assessment* 194(11): 833. <https://doi.org/10.1007/s10661-022-10479-9>.
- [38] Smith, M. D., Koerner, S. E., Knapp, A. K. (2020): Long-term response of grassland communities to climate variability. – *Ecology Letters* 23(4): 1028-1037.
- [39] Sutcliffe, M. E., Mountford, J. O., Smith, R. S. (2020): Seasonal shifts in plant diversity across European dry grasslands. – *Biodiversity and Conservation* 29(7): 2081-2101. <https://doi.org/10.1007/s10531-020-01974-1>.
- [40] Taherdoost, H. (2022): A Guide to Evaluate Academic Sources to Develop Research Paper: Source Selection in Academic Writing. – *Asian Review of Social Sciences* 11(1): 57-58. <https://doi.org/10.2139/ssrn.4178672>.
- [41] Wang, Y., Zhang, X., Li, J. (2021): Variations in Shannon–Wiener diversity in semi-arid plant communities. – *Journal of Vegetation Science* 32(2): e12949.

- [42] Zemmouri, B., Djebarni, M., Bouazza, M. (2021): Contribution of *Stipa tenacissima* and *Lygeum spartum* to soil stabilization and pastoral value in Algerian steppes. – *Journal of Arid Land Studies* 31(3): 97-104.
- [43] Zerrouki, M., Achour, H., Sahki, A. (2022): Resilience of steppe vegetation under climate variability and grazing pressure in central Algeria. – *Environmental Monitoring and Assessment* 194(2): 125. <https://doi.org/10.1007/s10661-021-09761-z>.
- [44] Zhang, Q., Li, Y., Yang, X. (2023): Climatic and human impacts on floristic composition in semi-arid ecosystems. – *Ecological Processes* 12(1): 10. <https://doi.org/10.1186/s13717-023-00406-2>.
- [45] Zhao, Y., Liu, Y., Chen, X. (2021): Effects of environmental pressures on plant diversity and evenness. – *Global Ecology and Conservation* 27: e01596.

APPENDIX

Appendix A. List of species and their flowering phenology (type and biological cycle)

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Centaurea pomeliana</i> (Batt.)	Asteraceae	Chamophyte	Spring-Summer	Biennial
<i>Myosotis versicolor</i> (Pers.) Smith = <i>M. discolor</i>	Boraginaceae	Thérophyte	Spring-Summer	Annual
<i>Nardurus maritimus</i> (L.) Janchen	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Silene tridentata</i> (Desf.)	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Adonis aestivalis</i> (L.)	Renonculaceae	Thérophyte	Spring-Summer	Annual
<i>Adonis dentata</i> Del.	Renonculaceae	Thérophyte	Spring-Summer	Annual
<i>Aegilops triuncialis</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Aeluropus littoralis</i> (Gouan.) Parl	Poaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Ajuga chamaepitys</i> (Schreber.)	Lamiaceae	Chamophyte	Spring-Summer	Annual
<i>Ajuga iva</i> (L.) Schreb	Lamiaceae	Chamophyte	Spring-Summer	Perennial
<i>Allium chamaemoly</i> (L.)	Liliaceae	Géophyte	Spring	Perennial
<i>Allium paniculatum</i> (L.) ssp. <i>typicum</i>	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Allium roseum</i> (L.)	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Alyssum alpestre</i> (L.)	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Alyssum granatense</i> Bois et Reut.	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Alyssum macrocalyx</i> Coss et Dur.	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Alyssum parviflora</i> Fisch.	Brassicaceae	Thérophyte	Spring-Summer	Annual
<i>Alyssum scutigerum</i> Dur.	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Ammodaucus leucotrichus</i> Coss. et Dur.	Apiaceae	Thérophyte	Spring-Summer	Annual
<i>Ammoides atlantica</i> (Coss. et Dur.) Wolf	Apiaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Ampelodesma mauritanica</i> (Poir.) Dur. et Schinz	Poaceae	Géophyte	Spring-Summer	Perennial

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Anacyclus clavatus</i> (Desf.)	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Anacyclus cyrtolepidioides</i> (Pomel.)	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Anacyclus radiatus</i> (Lois.)	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Anacyclus valentinus</i> (L.)	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Anarrhinum fruticosum</i> (Desf.)	Scrophulariaceae	Thérophyte	Spring-Summer	Perennial
<i>Androsace maxima</i> (L.)	Primulaceae	Thérophyte	Spring-Summer	Annual
<i>Anthemis pedunculata</i> Desf.	Asteraceae	Hémicryptophyte	Spring-Summer	Annual
<i>Anthyllis sericea</i> (L.) ssp. <i>sericea</i> (Lag.)	Fabaceae	Chaméphyte	Spring-Summer	Perennial
<i>Arabis auriculata</i> Lam.	Brassicaceae	Thérophyte	Spring-Summer	Annual
<i>Arabis verna</i> (L.) W.T. Aiton	Brassicaceae	Thérophyte	Spring-Summer	Annual
<i>Arceuthobium oxycedri</i> (DC.) M. Bieb.	Viscaceae	Hémiparasite	Spring-Summer	Perennial
<i>Arenaria serpyllifolia</i> (Rchb.) Guss	Caryophyllaceae	Thérophyte	Spring-Summer	Annual
<i>Aristida pungens</i> (Desf.)	Poaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Aristida ciliata</i> Desf.	Grammineae	Hémicryptophyte	Spring-Summer	Annual
<i>Artemisia campestris</i> (L.)	Asteraceae	Chamophyte	Spring-Summer	Perennial
<i>Artemisia herba-alba</i> (Asso)	Asteraceae	Chamophyte	Spring-Summer	Perennial
<i>Asparagus acutifolius</i> L.	Liliaceae	Phanérophyte	Spring-Summer	Perennial
<i>Astragalus armatus</i> Willd.	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus caprinus</i> (L.)	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus caprinus</i> ssp. <i>lanigerus</i> (Desf.) Maire	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus crucialis</i> (Link)	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus gombo</i> (Coss.) et Dur.	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus incanus</i> (L.)	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus monspessulanus</i> (L.)	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus scorioides</i> (Pourret)	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus senaicus</i> (Seniculus)	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Astragalus sesameus</i> (L.)	Fabaceae	Chamophyte	Spring-Summer	Perennial
<i>Atractylis echinata</i> (Pomel.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Atractylis caerulea</i> (Batt.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Atractylis cancellata</i> (L.) Steeppe	Asteraceae	Thérophyte	Spring-Summer	Perennial

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Atractylis carduus</i> (Forsk.) Christ.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Atractylis humilis</i> (L.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Atractylis humilis</i> ssp. <i>caespitosa</i> (Desf.) M.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Atractylis phaeolepis</i> (Pomel.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Atractylis polycephala</i> (Coss.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Atractylis serratuloides</i> (Sieb.)	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Atriplex halimus</i> (L.)	Chenopodiaceae	Nanophanérophte	Spring-Summer	Annual
<i>Avena breviaristata</i> (Barratte ex Trab.) Holub	Poaceae	Hémicryptophte	Spring-Summer	Annual
<i>Avena bromoides</i> (Gouan.)	Poaceae	Hémicryptophte	Spring-Summer	Annual
<i>Bascurtella didyma</i>	Crucifereae	Chamophte	Spring-Summer	Perennial
<i>Bassia muricata</i> (L.) Asch.	Chenopodiaceae	Thérophyte	Spring-Summer	Biennial
<i>Bellis silvestris</i> Cyrillo	Asteraceae	Hémicryptophte	Spring-Summer	Annual
<i>Beta vulgaris</i> (L.)	Chenopodiaceae	Thérophyte	Spring-Summer	Annual
<i>Biscutella didyma</i> (L.)	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Boreava aptera</i> Bois	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Brachypodium dichotomum</i> (L.) Maire (<i>Stoibrax dichotomum</i> (L.) Raf.)	Apiaceae	Thérophyte	Spring-Summer	Annual
<i>Bromus hordaceus</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Bromus madritensis</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Bromus ramosus</i> (L.) R. et S.	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Bromus rubens</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Bromus squarrosus</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Bromus tectorum</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Buffonia perennis</i> Pourr.	Caryophyllaceae/Alsinoideae	Chamophte	Spring-Summer	Perennial
<i>Buffonia tenuifolia</i> Griseb.	Caryophyllaceae/Alsinoideae	Chamophte	Spring-Summer	Perennial
<i>Bunium macuca</i> Boiss.	Apiaceae	Géophte	Spring-Summer	Perennial
<i>Bupleurum atlanticum</i> Murb.	Apiaceae	Hémicryptophte	Spring-Summer	Annual
<i>Bupleurum balansae</i> Boiss. et Reut.	Apiaceae	Hémicryptophte	Spring-Summer	Annual
<i>Calendula aegyptiaca</i> Desf.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Calendula arvensis</i> ssp. <i>communis</i> (Emb. et M.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Callitris articulata</i> (Vahl.) Link (<i>Thuja</i>)	Cupressaceae	Phanérophte	Spring-Summer	Perennial

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Calycotome spinosa</i> (L.) Lamk.	Fabaceae	PhanérophYTE	Spring-Autumn	Annual
<i>Calycotome villosa</i> (Poiret.) Link.	Fabaceae	PhanérophYTE	Spring-Summer	Perennial
<i>Capsella bursa-pastoris</i> (L.)	Crucifereae	Thérophyte	Spring-Summer	Perennial
<i>Carduncellus duvauxii</i> Batt.	Asteraceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Carduncellus pinnatus</i> (Desf.) DC.	Asteraceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Carduncellus plumosus</i> (Pomel.)	Asteraceae	Chamophyte	Spring-Summer	Perennial
<i>Carduncellus pomelianus</i> Batt.	Asteraceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Carduus getulus</i> Pomel.	Asteraceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Carex divisa</i> Huds.	Cyperaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Carex halleriana</i> Asso.	Cyperaceae	Hémicryptohyte	Spring-Summer	Annual
<i>Carlina involucrata</i> ssp. <i>corymbosa</i> (Q. et S.)	Asteraceae	Chaméphyte	Spring-Summer	Perennial
<i>Catapodium loliaceum</i> (Huds.) Link.	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Centaurea alba</i> L.	Asteraceae	Chamophyte	Spring-Summer	Perennial
<i>Centaurea melitensis</i> (L.)	Asteraceae	Chamophyte	Spring-Summer	Perennial
<i>Centaurea parviflora</i> (Desf.)	Asteraceae	Chamophyte	Spring	Annual
<i>Centaurea pengens</i> (Pomel)	Asteraceae	Chamophyte	Spring-Autumn	Annual
<i>Ceratocephalus falcatus</i> (L.) Pers. ssp. <i>incurvus</i> Stev.	Renonculaceae	Thérophyte	Spring-Autumn	Annual
<i>Chenopodium album</i> (L.)	Chenopodiaceae	Hémicryptohyte	Spring-Summer	Annual
<i>Chenopodium murale</i> (L.)	Chenopodiaceae	Hémicryptohyte	Spring-Summer	Annual
<i>Chrysanthemum fuscatum</i> Desf.	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Cistus villosus</i> L. var. <i>creticus</i>	Cistaceae	PhanérophYTE	Spring-Summer	Annual
<i>Cistus villosus</i> L. var. <i>mauritanicus</i> Grosser.	Cistaceae	PhanérophYTE	Spring-Summer	Annual
<i>Coronilla scorpioides</i> Koch.	Fabaceae	Thérophyte	Spring-Summer	Annual
<i>Ctenopsis pectinella</i> (Del.) De Not.	Poaceae	Thérophyte	Spring-Summer	Biennial
<i>Cutandia dichotoma</i> var. <i>memphitica</i> (Roth.) M. et W.	Poaceae	Thérophyte	Spring-Summer	Biennial
<i>Cutandia divaricata</i> (Desf.) Benth.	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Cutandia vallessina</i> (Honck.) Gaud.	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Cynoglossum cheirifolium</i> L.	Boraginaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Cynoglossum clandestinum</i> Desf.	Boraginaceae	Hémicryptohyte	Spring-Summer	Annual
<i>Cytisus fontanesii</i> Spach.	Fabaceae	Chaméphyte	Spring-Summer	Annual
<i>Dactylis glomerata</i> (L.)	Poaceae	Hémicryptohyte	Spring-	Annual

Species	Family	Biological Type	Flowering Period	Biological Cycle
		hyte	Summer	
<i>Delphinium pubescens</i> DC.	Renonculaceae	Thérophyte	Spring-Summer	Annual
<i>Diplotaxis eruroides</i> (L.)	Crucifereae	Thérophyte	Spring-Summer	Perennial
<i>Diplotaxis harra</i> (Forsk.) Boiss.	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Diplotaxis pitardiana</i> (Maire)	Crucifereae	Thérophyte	Spring-Summer	Perennial
<i>Diplotaxis virgata</i> DC.	Crucifereae	Thérophyte	Spring-Summer	Perennial
<i>Ebenus pinnata</i> L.	Fabaceae	Chaméphyte	Spring-Summer	Perennial
<i>Echinaria capitata</i> (L.) Desf.	Poaceae	Thérophyte	Summer	Perennial
<i>Echinops spinosus</i> L.	Asteraceae	Chamophyte	Autumn-Winter	Perennial
<i>Echium pycnanthum</i> (Pomel) ssp. <i>humile</i> (Desf.) Jah. et M.	Boraginaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Echium trygorrhizum</i> (Pomel.)	Boraginaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Epipactis helleborine</i> (L.) Crantz.	Orchidaceae	Géophyte	Spring-Summer	Annual
<i>Erica multiflora</i> (L.)	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Erodium bipinnatum</i> (Willd.)	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium botrys</i> (Cav.) Betel	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium cicutarium</i> (L'Her.)	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium glaucophyllum</i> (L'Her.)	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium hirtum</i> (Desf.)	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium hymenodes</i> (L'Her.)	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium malachoides</i> (L.) Willd.	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium meynieri</i> Maire.	Geraniaceae	Thérophyte	Spring-Summer	Annual
<i>Erodium polyanthemum</i> (Desf.) Pers.	Geraniaceae	Thérophyte	Spring-Summer	Perennial
<i>Erodium triangulare</i> ssp. <i>lacialum</i> (Cav.) Muschler	Geraniaceae	Thérophyte	Spring-Summer	Perennial
<i>Eruca vesicaria</i> (L.) Cav. ssp. <i>vesicaria</i> (L.) Briq.	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Eruca vesicaria</i> ssp. <i>pannatifida</i> (Desf.) Emb. et Maire	Brassicaceae	Thérophyte	Spring-Summer	Annual
<i>Eryngium ilicifolium</i> Lam.	Apiaceae	Thérophyte	Spring-Summer	Annual
<i>Erysimum bocconeii</i> (All.) Pers.	Brassicaceae	Thérophyte	Spring-Summer	Annual
<i>Erysimum incanum</i> Kunze	Brassicaceae	Thérophyte	Spring-Summer	Annual
<i>Euphorbia falcata</i> (L.)	Euphorbiaceae	Thérophyte	Spring-Summer	Annual
<i>Euphorbia chamaesyce</i> (L.)	Euphorbiaceae	Thérophyte	Spring-Summer	Annual
<i>Euphorbia exigua</i> (L.)	Euphorbiaceae	Thérophyte	Spring-Summer	Annual

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Euphorbia granulata</i> Forsk.	Euphorbiaceae	Thérophyte	Spring-Summer	Annual
<i>Euphorbia helioscopia</i> ssp. <i>helioscopiodes</i> (Losc. et Pardo.) Rouy	Euphorbiaceae	Thérophyte	Spring-Summer	Annual
<i>Euphorbia sulcata</i> de Lens.	Euphorbiaceae	Thérophyte	Spring-Summer	Perennial
<i>Evax argentea</i> Pomel.	Asteraceae	Thérophyte	Spring	Perennial
<i>Filago heterantha</i> (Guss.)	Asteraceae	Thérophyte	Spring	Perennial
<i>Filago spathulata</i> (Pers.)	Asteraceae	Thérophyte	Spring	Perennial
<i>Fumana thymifolia</i> (L.) Verlot.	Cistaceae	Chaméphyte	Spring	Perennial
<i>Gagea arvensis</i> (Pers.) Dumort.	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Gagea granatilli</i> (Parl.) ssp. <i>eu-granatilli</i> (Terracc.)	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Gagea mauritanica</i> Dur.	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Gagea reticulata</i> Pall.	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Galium palustre</i> L.	Scrophulariaceae	Chaméphyte	Spring-Summer	Annual
<i>Genista erioclada</i> ssp. <i>erioclada</i> (Emb.) et Maire.	Fabaceae	Chaméphyte	Spring-Summer	Annual
<i>Genista microcephala</i> (Coss.) et Dur.	Fabaceae	Chaméphyte	Spring	Perennial
<i>Genista quadriflora</i> Munby.	Fabaceae	Chaméphyte	Spring	Perennial
<i>Geranium dissectum</i> L.	Geraniaceae	Thérophyte	Spring	Perennial
<i>Geranium pyrenaicum</i> Burm.f.	Geraniaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Gladiolus byzantinus</i> Mill.	Iridaceae	Géophyte	Spring-Summer	Perennial
<i>Gladiolus byzantinus</i> Mill. (<i>G. communis byzantinus</i>)	Iridaceae	Géophyte	Spring-Summer	Perennial
<i>Gladiolus segetum</i> (Ker.) Gawl.	Iridaceae	Géophyte	Spring-Summer	Perennial
<i>Globularia alypum</i> L.	Globulariaceae	Chamaephyte	Spring-Summer	Perennial
<i>Helianthemum ledifolium</i> (L.) Mill.	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum cinereum</i> (Cav.) Pers. (= <i>Malcomia torulosa</i>)	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum hirtum</i> (Spreng.) ssp. <i>reficomum</i>	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum hirtum</i> ssp. <i>ruficomum</i> (Viv.) M.	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum kahiricum</i>	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum lipii</i> (L.) Pers.	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum papillare</i> (Boiss.)	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum pilosum</i> (L.) Pers.	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum virgatum</i>	Cistaceae	Chamophyte	Spring-Summer	Perennial
<i>Helianthemum apertum</i> (Pomel.)	Cistaceae	Chamophyte	Spring-Summer	Annual
<i>Helichrysum stoechas</i> (L.) DC.	Asteraceae	Chaméphyte	Spring-Summer	Annual
<i>Helosciadium nodiflorum</i> Lag.	Apiaceae	Hémicryptop	Spring-	Annual

Species	Family	Biological Type	Flowering Period	Biological Cycle
		hyte	Summer	
<i>Herniaria fontanesii</i> J. (=fruticosa (L.) ssp.)	Caryophyllaceae	Thérophyte	Spring-Summer	Annual
<i>Herniaria glabra</i> L.	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Herniaria hirsuta</i> L.	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Herniaria incana</i> Lamk.	Caryophyllaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Herniaria mauritanica</i> (Murb.) Fontanisii Batt.	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Hertia cheirifolia</i> (L.) O.K.	Poaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Hippocrepis bicontorta</i> (Loisel.)	Fabaceae	Thérophyte	Spring	Annual
<i>Hippocrepis multisiliquosa</i> (L.) ssp. <i>ciliata</i> (Willd.) Maire	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Hippocrepis unisiliquosa</i> (L.)	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Hordeum murinum</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Annual
<i>Hypocoum pendulum</i> (L.)	Papaveraceae	Thérophyte	Spring-Summer	Annual
<i>Hypochoeris laevigata</i>	Asteraceae	Hémicryptophyte	Spring	Perennial
<i>Hypochoeris radicata</i> L.	Asteraceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Iberis odorata</i> L.	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Ifloga spicata</i> (Forsk.) Sch. Bip.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Iris sisyrinchium</i> L.	Iridaceae	Géophyte	Spring-Summer	Perennial
<i>Juniperus oxycedrus</i> L.	Cupressaceae	Phanérophyte	Spring-Summer	Perennial
<i>Juniperus phoenicea</i> L.	Cupressaceae	Phanérophyte	Spring-Summer	Annual
<i>Jurinea humilis</i> DC.	Asteraceae	Hémicryptophyte	Spring-Summer	Annual
<i>Knautia arvensis</i> (L.) Coult.	Dipsacaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Koeleria pubescens</i> (Lank.) P.B.	Poaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Koelpinia linearis</i> Pallas	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Koniga lybica</i> (Viv.)	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Lagurus ovatus</i> L.	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Lappula redowskii</i> (Hornem.) Greene	Boraginaceae	Thérophyte	Spring-Summer	Perennial
<i>Launaea acanthoclada</i> M.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Launaea angustifolia</i> (Desf.) Muschler	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Launaea arborescens</i> (Batt.) M.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Launaea glomerata</i> (Cass.) Hook.f.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Launaea resedifolia</i> ssp. <i>eu-residifolia</i> (M.)	Asteraceae	Thérophyte	Spring-Summer	Perennial

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Leontodon hispanicus</i> (Del.) Boiss.	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Leontodon hispidulus</i> (Del.) Boiss.	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Leontodon saxatilis</i> (Lamk.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Leontodon tuberosus</i> (L.)	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Lepidium subulatum</i> L.	Asteraceae	Thérophyte	Spring	Perennial
<i>Lepturus cylindricus</i> (Willd.) Trin.	Crucifereae	Chaméphyte	Spring	Perennial
<i>Leuzea conifera</i> (L.) DC.	Poaceae	Thérophyte	Spring	Perennial
<i>Linaria simplex</i> (DC.)	Asteraceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Linum gallicum</i> (L.)	Scrophulariaceae	Chamophyte	Spring-Summer	Perennial
<i>Linum decumbens</i> Desf.	Linaceae	Hémiryphtophyte	Spring	Annual
<i>Linum strictum</i> (L.)	Linaceae	Hémiryphtophyte	Spring-Summer	Perennial
<i>Linum suffruticosum</i> (L.)	Linaceae	Hémiryphtophyte	Spring-Summer	Perennial
<i>Lithospermum apulum</i> (L.) Vahl.	Linaceae	Hémiryphtophyte	Spring	Annual
<i>Lithospermum arvense</i> (L.)	Boraginaceae	Thérophyte	Spring-Summer	Annual
<i>Lobularia lybica</i> (Viv.) Webb	Boraginaceae	Thérophyte	Spring-Summer	Perennial
<i>Lobularia maritima</i> (L.) Desv.	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Loeflingia hispanica</i> L.	Crucifereae	Hémicryptophyte	Spring-Summer	Perennial
<i>Lolium multiflorum</i> Lam.	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Lolium perenne</i> (L.)	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Lolium rigidum</i> (Gaud.)	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Lonicera implexa</i> L.	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Lotus creticus</i> (L.) ssp. <i>collinus</i> (Boiss.) Briquet	Caprifoliaceae	Phanérophyte	Spring-Summer	Perennial
<i>Lotus creticus</i> (L.) ssp. <i>eu-creticus</i> (Briquet.)	Fabaceae	Chaméphyte	Spring-Summer	Perennial
<i>Lotus edulis</i> L.	Fabaceae	Chaméphyte	Spring-Summer	Perennial
<i>Lotus jolyi</i> Batt.	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Lotus ornithopodioides</i> L.	Fabaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Lotus parviflorus</i> Desf.	Fabaceae	Thérophyte	Spring-Summer	Annual
<i>Lotus pusillus</i> Medik.	Fabaceae	Thérophyte	Spring-Summer	Annual
<i>Lotus villosus</i> (Forsk.) <i>pusillus</i> (Medik.)	Fabaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Lygeum spartum</i> (L.)	Fabaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Malcolmia aegyptiaca</i> Spr.	Poaceae	Hémicryptophyte	Spring-Summer	Annual

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Malcolmia torulosa</i> (Desf.) Boiss.	Crucifereae	Thérophyte	Spring-Summer	Perennial
<i>Malva aegyptiaca</i> (L.)	Crucifereae	Thérophyte	Spring	Annual
<i>Malva aegyptiaca</i> (L.)	Malvaceae	Thérophyte	Spring-Summer	Perennial
<i>Malva parviflora</i> (L.)	Malvaceae	Thérophyte	Spring-Summer	Perennial
<i>Malva sylvestris</i> (L.)	Malvaceae	Thérophyte	Spring-Summer	Perennial
<i>Mantisalca salmantica</i> (L.) Brq et Cavill	Malvaceae	Thérophyte	Spring-Summer	Perennial
<i>Matthiola fruticosa</i> (L.) Maire	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Matthiola longipetala</i> (Vent.) DC. ssp. <i>livida</i> (Del.) Maire	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Matthiola longipetala</i> ssp. <i>kralikii</i> (Pomel.) Maire	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Medicago italica</i> (Mill.) Steud. ssp. <i>helix</i> (Willd.) Emb. et Maire	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Medicago laciniata</i> (L.) All.	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Medicago litoralis</i> (Rohde.)	Fabaceae	Thérophyte	Spring-Summer	Annual
<i>Medicago minima</i> (Gruf.)	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Medicago truncatula</i> (Gearn.)	Fabaceae	Thérophyte	Spring	Perennial
<i>Medicago turbinata</i> (L.) Wild.	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Melilotus indica</i> (L.) All	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Mentha pulegium</i> L.	Fabaceae	Thérophyte	Spring-Summer	Annual
<i>Micropus bombicinus</i> (Lag.)	Lamiaceae	Chaméphyte	Spring-Summer	Perennial
<i>Minuartia campestris</i> (L.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Minuartia tenuifolia</i> (L.) Hiern.	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Moricandia arvensis</i> (L.) DC.	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Mpelodesma mauritanica</i>	Brassicaceae	Chamophyte	Spring-Summer	Perennial
<i>Muricaria prostrata</i> (Desf.)	Plantaginaceae	Thérophyte	Spring-Summer	Perennial
<i>Muscari comosum</i> (L.) Mill.	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Muscari maritimum</i> Desf.	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Muscari parviflora</i> Desf.	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Myosotis scorpioides</i> var. <i>collina</i> Ehrh	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Myosotis scorpioides</i> var. <i>collina</i> Ehrh	Boraginaceae	Thérophyte	Spring-Summer	Perennial
<i>Nardurus cynosuroides</i> (Desf.) B. et T.	Boraginaceae	Thérophyte	Spring-Summer	Perennial
<i>Noaea mucronata</i> (Forsk.) Asch.	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Nonnea micronatha</i> Boiss. et Reut.	Chenopodiaceae	Chamophyte	Summer	Perennial

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Nonnea visicaria</i> (L.) Rsch.	Boraginaceae	Thérophyte	Summer	Perennial
<i>Olea sylvestris</i> L.	Boraginaceae	Thérophyte	Spring-Summer	Perennial
<i>Onobrychis viciifolia</i> Scop.	Oleaceae	Phanérophyte	Spring	Perennial
<i>Onopordon acaule</i> (L.)	Fabaceae	Hémicryptohyte	Spring-Summer	Annual
<i>Onopordon arenarium</i> (Desf) Pomel	Asteraceae	Hémicryptohyte	Spring-Summer	Annual
<i>Ophrys lutea</i> (Cav.) Gouan	Asteraceae	Hémicryptohyte	Spring-Summer	Annual
<i>Orchis</i> sp. L.	Orchidaceae	Géophyte	Spring-Summer	Perennial
<i>Papaver rhoeas</i> (L.)	Orchidaceae	Géophyte	Spring-Summer	Perennial
<i>Papaver hybridum</i> (L.)	Papaveraceae	Thérophyte	Spring-Summer	Annual
<i>Papaver malviflorum</i> (Droum.)	Papaveraceae	Thérophyte	Spring-Summer	Perennial
<i>Paronychia capitata</i> ssp. <i>chlorothyrum</i> (Murberk.) Maire	Papaveraceae	Thérophyte	Spring-Summer	Perennial
<i>Paronychia arabica</i> (L.) Dc ssp. <i>cossoniana</i> (Gay)	Caryophyllaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Paronychia arabica</i> ssp. <i>annua</i> (Del) Maire	Caryophyllaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Paronychia argentea</i> (Pourr.) Lamk	Caryophyllaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Paronychia capitata</i> (L.) Lamk	Caryophyllaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Peganum harmala</i> (L.)	Caryophyllaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Phagnalon rupestre</i> (L.) DC.	Zygophyllaceae	Chamophyte	Spring-Summer	Perennial
<i>Phagnalon saxatile</i> (L.) Cass.	Asteraceae	Chamophyte	Spring-Summer	Perennial
<i>Phillyrea angustifolia</i> L.	Asteraceae	Chamophyte	Spring-Summer	Annual
<i>Pinus halepensis</i> Mill.	Oleaceae	Phanérophyte	Spring-Summer	Annual
<i>Pistacia lentiscus</i> (L.)	Pinaceae	Phanérophyte	Spring-Summer	Annual
<i>Pistacia terebenthus</i> (L.)	Anacardiaceae	Hémicryptohyte	Spring-Summer	Annual
<i>Plantago albicans</i> L.	Anacardiaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Plantago amplexicaule</i> Cav.	Plantaginaceae	Hémicryptohyte	Summer	Annual
<i>Plantago ciliata</i>	Plantaginaceae	Thérophyte	Summer	Annual
<i>Plantago psyllium</i> ((L.))	Plantaginaceae	Thérophyte	Spring-Summer	Perennial
<i>Poa bulbosa</i> (L.)	Plantaginaceae	Thérophyte	Spring-Summer	Perennial
<i>Polygonum aviculare</i> (L.)	Poaceae	Hémicryptohyte	Spring-Summer	Perennial
<i>Polygonum equisetiforme</i> S et Sm.	Polygonaceae	Thérophyte	Spring-Summer	Annual
<i>Pseudognaphalium luteoalbum</i> (L.) Hilliard & B.L.Burt	Polygonaceae	Chaméphyte	Spring-Summer	Annual
<i>Pulicaria laciniata</i> (Coss et Gral.) Thel	Asteraceae	Thérophyte	Spring-Summer	Perennial

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Quercus ilex L.</i>	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Ranunculus arvensis (L)</i>	Fagaceae	Phanérophyte	Spring-Summer	Annual
<i>Ranunculus gramineus L.</i>	Renonculaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Reseda alba (L.) ssp. eu-alba Maire</i>	Renonculaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Reseda arabica (Boiss)</i>	Resedaceae	Thérophyte	Spring-Summer	Perennial
<i>Reseda decursiva (Forsk.)</i>	Resedaceae	Thérophyte	Spring-Summer	Perennial
<i>Reseda sp. (Duriaeana) L.</i>	Resedaceae	Thérophyte	Spring-Summer	Perennial
<i>Roemeria hybrida (L.) DC.</i>	Resedaceae	Thérophyte	Spring-Summer	Perennial
<i>Romulea bolbucodium (L.) Seb.</i>	Papaveraceae	Thérophyte	Spring-Summer	Perennial
<i>Rosmarinus officinalis L.</i>	Iridaceae	Géophyte	Spring-Summer	Annual
<i>Rosmarinus tournifortii De Noé</i>	Lamiaceae	Phanérophyte	Spring-Summer	Perennial
<i>Rubia peregrina L.</i>	Lamiaceae	Phanérophyte	Spring-Summer	Perennial
<i>Rumex gonglomeratus (Murr.) (L'SSAN ELFARD)</i>	Rubiaceae	Phanérophyte	Spring-Summer	Perennial
<i>Rumex aristidis Coss.</i>	Polygonaceae	Thérophyte	Spring-Summer	Perennial
<i>Saccocalyx satureioides (Coss.) et Dur</i>	Polygonaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Salicornia arabica L. var. perennis (Mill.) Fiori</i>	Lamiaceae	Chaméphyte	Spring-Summer	Perennial
<i>Salicornia arabica L. var. typica Fiori</i>	Chenopodiaceae	Chaméphyte	Spring-Summer	Perennial
<i>Salsola foetida (Del.)</i>	Chenopodiaceae	Chaméphyte	Summer	Annual
<i>Salsola vermiculata (L.)</i>	Chenopodiaceae	Chamophyte	Summer	Annual
<i>Salvia verbenaca ssp. clandestina (L.) Pugsl</i>	Chenopodiaceae	Chamophyte	Spring-Summer	Annual
<i>Scabiosa arenarea Forsk.</i>	Lamiaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Scabiosa atropurpurea L.</i>	Dipsacaceae	Thérophyte	Spring-Summer	Perennial
<i>Scabiosa stellata L.</i>	Dipsacaceae	Thérophyte	Spring-Summer	Perennial
<i>Schismus barbatus (L.) (Thell)</i>	Dipsacaceae	Thérophyte	Spring-Summer	Perennial
<i>Scilla lingulata Poiret.</i>	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Scilla obtusifolia Poiret.</i>	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Scilla peruviana L.</i>	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Scorzonera laciniata (L.)</i>	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Scorzonera undelata ssp. alexandria (Boiss.) M</i>	Asteraceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Scrofularia laevigata Vahl.</i>	Asteraceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Sedum sediforme (Jacq.) Pau.</i>	Scrophulariaceae	Hémicryptophyte	Spring-Summer	Annual

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Senecio flavus</i> (Dec.) Sch.Bip.	Crassulaceae	Thérophyte	Spring-Summer	Perennial
<i>Senecio gallicus</i> ssp. <i>coronopifolius</i> (Desf.) M	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Senecio leucanthemifolius</i> Poir.	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Senecio vulgaris</i> ssp. <i>massaicus</i> (Maire.) Q.et S.	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Sideritis montana</i> (L.) subsp.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Silene arenarioides</i> (Desf.)	Lamiaceae	Thérophyte	Spring-Summer	Annual
<i>Silene cerastioides</i> (L.)	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Silene conica</i> (L.)	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Silene rouyana</i> (B.)	Caryophyllaceae	Thérophyte	Spring-Summer	Annual
<i>Silene rubella</i> (L.)	Caryophyllaceae	Thérophyte	Spring-Summer	Annual
<i>Silene secundiflora</i> Otth.	Caryophyllaceae	Thérophyte	Spring-Summer	Annual
<i>Silene setacea</i> (Viv.)	Caryophyllaceae	Thérophyte	Spring-Summer	Annual
<i>Sisymbrium irio</i> (L.)	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Sisymbrium runcinatum</i> Lag.	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Sisymbrium thalianum</i> (L.) Gay. et Mon. (= <i>Arabidopsis thaliana</i> (L.) Heyn.)	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Sisymbrium torulosum</i> (Desf.)	Crucifereae	Thérophyte	Spring-Summer	Annual
<i>Smilax aspera</i> L.	Crucifereae	Thérophyte	Spring-Summer	Perennial
<i>Sonchus oleraceus</i> L.	Lamiaceae	Nanophanérophte	Spring-Summer	Perennial
<i>Spergula arvensis</i> L.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Spergula flaccida</i> (Roxb.) Asch	Caryophyllaceae	Thérophyte	Spring-Summer	Perennial
<i>Sphenopus divaricatus</i> (Gouan.) Rchb.	Caryophyllaceae	Hémicryptophte	Spring-Summer	Perennial
<i>Spitzelia coronopifolia</i> Desf.	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Leontodon hispanicus</i> (Del.) Boiss.	Asteraceae	Hémicryptophte	Spring-Summer	Perennial
<i>Stachys brachyclada</i> Noé ex Coss	Lamiaceae	Thérophyte	Spring-Summer	Perennial
<i>Stachys circinata</i> L'Her.	Lamiaceae	Hémicryptophte	Spring-Summer	Perennial
<i>Stahelina dubia</i> (L.)	Asteraceae	Chaméphyte	Spring-Summer	Perennial
<i>Stipa barbata</i> (Desf.)	Poaceae	Hémicryptophte	Spring-Summer	Perennial
<i>Stipa parviflora</i> (Desf.)	Poaceae	Hémicryptophte	Spring-Summer	Perennial
<i>Stipa retorta</i> (Cav.)	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Stipa tenacissima</i> (L.)	Poaceae	Hémicryptophte	Spring-Summer	Perennial

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Suaeda fruticosa</i> L.	Chenopodiaceae	Chaméphyte	Spring-Summer	Perennial
<i>Tamarix aphylla</i> (L.) Karst	Tamaricaceae	Phanérophyte	Spring-Summer	Perennial
<i>Tamarix gallica</i> L.	Tamaricaceae	Phanérophyte	Spring-Summer	Perennial
<i>Teledium imperati</i> (L.)	Caryophyllaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Teledium spaerospermum</i> Boiss.	Caryophyllaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Tetraclinis articulata</i> (Vahl.) Link (= <i>Callitris articulata</i> (Vahl.) Link)	Cupressaceae	Phanérophyte	Spring-Summer	Perennial
<i>Tetraclinis articulata</i> (Vahl.) Link Masters. <i>Thuja de Barbarie</i>	Cupressaceae	Phanérophyte	Spring-Summer	Annual
<i>Teucrium polium</i> (L.) p.p.	Lamiaceae	Chamophyte	Spring-Summer	Perennial
<i>Teucrium pseudo-chamaepitys</i> (L.)	Lamiaceae	Chamophyte	Spring-Summer	Perennial
<i>Thapsia vilosa</i> L.	Apiaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Thlaspi perfoliatum</i> L.	Brassicaceae	Thérophyte	Spring-Summer	Perennial
<i>Thymelaea tartonraira</i> All.	Theligonaceae	Chamophyte	Spring-Summer	Perennial
<i>Thymus algeriensis</i> (Boiss.) et Reut	Lamiaceae	Chamophyte	Spring-Summer	Annual
<i>Thymus ciliatus</i> (Desf.) ssp. <i>eu-ciliatus</i> (Maire.)	Lamiaceae	Chamophyte	Spring-Summer	Annual
<i>Thymus hirtus</i> Willd.	Lamiaceae	Chamophyte	Spring-Summer	Annual
<i>Tourneuxia variifolia</i> (Coss.)	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Tragopogon porrifolius</i> L.	Asteraceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Trigonella monspeliaca</i> (L.)	Fabaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Trigonella polycerata</i> L.	Fabaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Trisetaria parviflora</i> (Desf.) Maire	Poaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Tuberaria guttata</i> (L.) Foureau	Cistaceae	Thérophyte	Spring-Summer	Perennial
<i>Tulipa silvestris</i> (L.) ssp. <i>Australis</i> (Link.) Pamp	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Tulipa silvestris</i> (L.) ssp. <i>primulina</i> (Beker.) M.W	Liliaceae	Géophyte	Spring-Summer	Perennial
<i>Tulipa silvestris</i> ssp. <i>australis</i> (Link.) Pamp.	Liliaceae	Géophyte	Spring-Summer	Annual
<i>Tunica illyrica</i> (Ard.) Fisch et Meg (= <i>Dianthella compressa</i> (Claus.))	Caryophyllaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Tunica illyrica</i> (Ard.) Fisch et Meg.	Caryophyllaceae	Hémicryptophyte	Spring-Summer	Annual
<i>Turgenia latifolia</i> L. (Hoffm.)	Apiaceae	Thérophyte	Spring-Summer	Annual
<i>Urospermum picroides</i> (L.) Schmidt	Asteraceae	Thérophyte	Spring-Summer	Annual
<i>Valeriana tuberosa</i> L.	Valerianaceae	Hémicryptophyte	Spring-Summer	Perennial
<i>Valerianella carinata</i> Lois.	Valerianaceae	Thérophyte	Spring-Summer	Annual

Species	Family	Biological Type	Flowering Period	Biological Cycle
<i>Valerianella coronata</i> (L.) DC.	Valerianaceae	Thérophyte	Spring-Summer	Annual
<i>Verbena supina</i> L.	Verbenaceae	Thérophyte	Spring-Summer	Annual
<i>Veronica praecox</i> All.	Scrophulariaceae	Thérophyte	Spring-Summer	Annual
<i>Vicia ervilia</i> (L.) Willd.	Fabaceae	Thérophyte	Spring-Summer	Annual
<i>Vicia monardi</i> (Boiss.)	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Vicia sativa</i> (L.)	Fabaceae	Thérophyte	Spring-Summer	Perennial
<i>Vulpia membranacea</i> (L.) Link	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Vulpia myuros</i> (L.) Gmel.	Poaceae	Thérophyte	Spring-Summer	Perennial
<i>Xeranthemum inapertum</i> (L.) Mill.	Asteraceae	Thérophyte	Spring-Summer	Perennial
<i>Ziziphora hispanica</i> (L.)	Lamiaceae	Thérophyte	Spring-Summer	Perennial
<i>Ziziphora officinalis</i> (L.)	Lamiaceae	Thérophyte	Spring-Summer	Perennial
<i>Ziziphus lotus</i>	Rhamnaceae	Phanerophyte	Spring-Summer	Perennial
<i>Zizyphora hispanica</i> L.	Lamiaceae	Thérophyte	Spring-Summer	Perennial